

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 September 2004 (02.09.2004)

PCT

(10) International Publication Number
WO 2004/073656 A2

- (51) International Patent Classification⁷: A61K (74) Agents: SANDBAKEN, Mark, G. et al.; TOWNSEND and TOWNSEND and CREW LLP, Two Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).
- (21) International Application Number: PCT/US2004/005247
- (22) International Filing Date: 20 February 2004 (20.02.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/449,055 20 February 2003 (20.02.2003) US
- (71) Applicant (for all designated States except US): SEATTLE GENETICS, INC. [US/US]; 21823 - 30th Drive, S.E., Bothell, WA 98021 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): LAW, Che-Leung [US/US]; 18834 Fremont Avenue N., Shoreline, WA 98133 (US). WAHL, Alan, F. [US/US]; 6150 East Mercer Way, Mercer Island, WA 98020 (US). SCHOLLER, Nathalie [FR/US]; 6208 32nd Avenue N.W., Seattle, WA 98107 (US). PESTANO, Linda, A. [US/US]; 11630 North Dragoon Springs Drive, Oro Valley, AZ 85737 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: ANTI-CD70 ANTIBODY-DRUG CONJUGATES AND THEIR USE FOR THE TREATMENT OF CANCER AND IMMUNE DISORDERS

IF6 VL nucleotide sequence:

```

1 atggagacag acacactcct gttatgggta ctgtgctctt gggtccagg ttcactggt 60
61 gacatgtgtc tgacacagtc tccgtcttcc ttacgtgtat ctctgggcca gagggccacc 120
121 atctcatgca gggccagcaa aggtgtcagt acatctggct atagtttat gcaactgtat 180
181 caacagaaac caggccagcc acccaaacct ctcctctata ttgtatcaa cctagaatct 240
241 ggggtccctg ccaggttcag tggcagtggt tctgggacag acttcaacct caacatcoat 300
301 cctgtggagg aggaggtgto tgcaacctat tactgtcagc acagtatgga ggttccgtgg 360
361 acgttcgggt gaggtaccac gctggaaato aaacgg 396

```

IF6 VL amino acid sequence:

```

-20 METDTLLLV LLLVFGSTC 0
1 DIVLTSPAS LAYSLGURAT ISCRASKSVS TSGVSPHRY QDEPGIRPKL LIYLASHLES 60
61 GVPARFSGSG SGDTETLRIH FVEREDATFY VOHSREVPW TRGGGTLEI KR 112

```

IF6 VH nucleotide sequence:

```

1 atggcttggg ttggacctt gctattctgt atggcagctg cccaaagtgc ccaagcacag 60
61 atccagtggg tgcagtcttg acctaggttg aagaagcttg gagggacagt caagatctcc 120
121 tgcagggttt ctgggtatc cttcaaaac tatggaatga actgggtgaa gcaggctcca 180
181 ggaagggttt taagtggat gggctggata aacacctaca ctggagagcg aacatatgct 240
241 gatgcttcca agggaggttt tgccttctct ttggaacct ctgccagcac tgcctatttg 300
301 cagatcaaca acctcaaaa tgggagacag gctacatatt tctgtgaaag agactacggc 360
361 gactatgta ttgactactg ggttcaagga aactcagta cgtctctctc a 411

```

IF6 VH amino acid sequence:

```

-19 HALVTHLPL EAARQSGA 0
1 QIDLVDSGE VKKPGTVMV SCASGTYFT EYCHMAMIA PGGLAMGN EHTTGEFTY 60
61 ADAFGKRFAP SLETSASTAY LQINSLKED TATYFCARDY GDIGDYNGU QTSVTVSS 118

```

(57) Abstract: Disclosed are anti-CD70 antibodies and derivatives thereof conjugated to cytotoxic, immunosuppressive, or other therapeutic agents, as well as pharmaceutical compositions and kits comprising the antibody- and antibody derivative-drug conjugates. Also disclosed are methods, for the treatment of CD70-expressing cancers and immunological disorders, comprising administering to a subject the disclosed pharmaceutical compositions.

WO 2004/073656 A2

BEST AVAILABLE COPY



Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5 **ANTI-CD70 ANTIBODY-DRUG CONJUGATES AND THEIR USE FOR
THE TREATMENT OF CANCER AND IMMUNE DISORDERS**

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application No.
10 60/449,055, filed Feb. 20, 2003, which is incorporated by reference herein in its entirety.

BACKGROUND OF THE INVENTION

[0002] CD70 is a member of the tumor necrosis factor (TNF) family of cell membrane-bound and secreted molecules that are expressed by a variety of normal and malignant cell types.
15 The primary amino acid (AA) sequence of CD70 predicts a transmembrane type II protein with its carboxyl terminus exposed to the outside of cells and its amino terminus found in the cytosolic side of the plasma membrane (Bowman *et al.*, 1994, *J Immunol* 152:1756-61; Goodwin *et al.*, 1993, *Cell* 73:447-56). Human CD70 is composed of a 20 AA cytoplasmic domain, an 18 AA transmembrane domain, and a 155 AA extracytoplasmic domain with two
20 potential N-linked glycosylation sites (Bowman *et al.*, *supra*; Goodwin *et al.*, *supra*). Specific immunoprecipitation of radioisotope-labeled CD70-expressing cells by anti-CD70 antibodies yields polypeptides of 29 and 50 kDa (Goodwin *et al.*, *supra*; Hintzen *et al.*, 1994, *J Immunol* 152:1762-73). Based on its homology to TNF-alpha and TNF-beta, especially in structural strands C, D, H and I, a trimeric structure is predicted for CD70 (Petsch *et al.*,
25 1995, *Mol Immunol* 32:761-72).

[0003] Original immunohistological studies revealed that CD70 is expressed on germinal center B cells and rare T cells in tonsils, skin, and gut (Hintzen *et al.*, 1994, *Int Immunol* 6:477-80). Subsequently, CD70 was reported to be expressed on the cell surface of recently antigen-activated T and B lymphocytes, and its expression wanes after the removal of
30 antigenic stimulation (Lens *et al.*, 1996, *Eur J Immunol* 26:2964-71; Lens *et al.*, 1997, *Immunology* 90:38-45). Within the lymphoid system, natural killer cells (Orengo *et al.*, 1997, *Clin Exp Immunol* 107:608-13) and mouse mature peripheral dendritic cells (Akiba *et al.*, 2000, *J Exp Med* 191:375-80) also express CD70. In non-lymphoid lineages, CD70 has been detected on thymic medullar epithelial cells (Hintzen *et al.*, 1994, *supra*; Hishima *et al.*,
35 2000, *Am J Surg Pathol* 24:742-46).

- 5 [0004] In addition to expression on normal cells, CD70 expression has been reported in different types of cancers including lymphomas, carcinomas, and tumors of neural origin. In malignant B cells, 71% of diffuse large B-cell lymphomas, 33% of follicle center lymphomas, 25% of mantle lymphomas, and 50% of B-CLL have been reported to express CD70 (Lens *et al.*, 1999, *Br J Haematol* 106:491-503). CD70 is frequently expressed
10 together with other lymphoid activation markers on the malignant Hodgkin and Reed-Sternberg cells of Hodgkin's disease (Gruss and Kadin, 1996, *Baillieres Clin Haematol* 9:417-46). One report demonstrates CD70 expression on 88% (7 of 8 cases) of thymic carcinomas and 20% (1 of 5 cases) of atypical thymomas (Hishima *et al.*, 2000, *supra*). The second type of carcinoma on which CD70 has been detected is nasopharyngeal carcinoma. One study
15 reports the presence of CD70 on 80% (16 of 20 cases) of snap-frozen tumor biopsies obtained from undifferentiated nasopharyngeal carcinomas (Agathangelou *et al.*, 1995, *Am J Path* 147:1152-60). CD70 has also been detected on brain tumor cells, especially glioma cell lines, solid human gliomas, and meningiomas (Held-Feindt and Mentlein, 2002, *Int J Cancer* 98:352-56; Wischusen *et al.*, 2002, *Can Res* 62:2592-99).
- 20 [0005] It has been proposed that transforming viruses including the Epstein-Barr virus (EBV) and the human T leukemia virus-1 (HTLV-1) can induce CD70 on cells such as epithelial cells that normally do not express CD70 (Agathangelou *et al.*, *supra*; Stein *et al.*, 1989, Oxford University Press, p. 446). Therefore, expression of CD70 on malignant B cells could be a reflection of oncogenic transformation (Lens *et al.*, 1999, *supra*). Also, since CD70
25 expression is induced on B cells after antigen encounter (Maurer *et al.*, 1990, *Eur J Immunol* 20:2679-84; Lens *et al.*, 1996, *supra*), stable expression of CD70 might reflect prolonged antigenic stimulation. This has been postulated to occur in follicular non-Hodgkin's lymphomas based on ongoing somatic hypermutation (Bahler *et al.*, 1992, *Proc Natl Acad Sci USA* 89:6770-74; Bahler *et al.*, 1992, *Cancer Res* 52:suppl. 5547S-51S).
- 30 [0006] The receptor for CD70 is CD27, a glycosylated type I transmembrane protein of about 55kDa (Goodwin *et al.*, 1993, *Cell* 73:447-56; Hintzen *et al.*, 1994, *supra*). CD70 is sometimes referred to as CD27L. CD27, which exists as a homodimer on the cell surface (Gravestien *et al.*, 1993, *Eur J Immunol* 23:943-50), is a member of the TNF receptor
35 superfamily as defined by cysteine-rich repeats of about 40 amino acids in the extracellular domain (Smith *et al.*, 1990, *Science* 248:1019-23; Locksley *et al.*, 2001, *Cell* 104:487-501). CD27 is typically expressed by thymocytes, NK, T, and B cells (Hintzen *et al.*, 1994, *Immunol Today* 15:307-11; Lens *et al.*, 1998, *Semin Immunol* 10:491-99). On resting T cells,

5 CD27 is constitutively expressed, yet antigenic triggering further upregulates CD27 expression (de Jong *et al.*, 1991, *J Immunol* 146:2488-94; Hintzen *et al.*, 1993, *J Immunol* 151:2426-35). Further, triggering of T cells via their T cell antigen receptor complex alone or in combination with the accessory molecule CD28 releases soluble CD27 from activated T cells (Hintzen *et al.*, 1991, *J Immunol* 147:29-35). Naïve B cells do not express CD27, but its
10 expression is induced and, in contrast to CD70, sustained after antigenic triggering of B cells (Jacquot S *et al.*, 1997 *J Immunol* 159:2652-57; Kobata T *et al.*, 1995, *Proc Natl Acad Sci USA* 92:11249-53).

[0007] In marked contrast to the restricted expression of CD27 and CD70 in normal B lineage cells, both CD27 and CD70 are frequently co-expressed in many B cell non-
15 Hodgkin's lymphomas and leukemias. This could potentially lead to functional CD27-CD70 interactions on these cells in the form of an autocrine loop, resulting in CD27 signaling and in CD70-induced proliferation, thereby providing a growth advantage to malignant cells (Lens *et al.*, 1999, *supra*).

[0008] The available data supports a model in which ligation of CD27 on activated
20 lymphocytes by CD70 delivers signals to the CD27-expressing cells, including co-stimulatory signals in T, B, and NK cells. (See, e.g., Goodwin *et al.*, *supra*; Hintzen *et al.*, 1995, *J Immunol* 154:2612-23; Oshima *et al.*, 1998, *Int Immunol* 10:517-26; Smith *et al.*, *supra*; Van Lier *et al.*, 1987, *J Immunol* 139:1589-96; Gravestein *et al.*, 1995, *Int Immunol* 7:551-7; Tesselaar *et al.*, 1997, *J Immunol* 159:4959-65; Jacquot *et al.*, *supra*; Agematsu *et al.*, 1998, *Blood* 91:173-80; Kobata *et al.*, *supra*; Agematsu *et al.*, 1997, *Eur J Immunol* 27:2073-79; Sugita *et al.*, 1992, *J Immunol* 149:1199-1203; Orengo *et al.*, 1997, *Clin Exp Immunol* 107:608-13). Antibodies against both murine and human CD70 have been demonstrated to inhibit such activities, presumably by blocking the CD70/CD27 interaction (Hintzen *et al.*, 1994, *supra*; Hintzen *et al.*, 1995, *supra*; Oshima *et al.*, *supra*).

30 [0009] Limited information is available on the modulation of cellular functions through CD70 signaling upon CD70/CD27 interaction, i.e., 'reverse signaling'. Some CD70 antibodies have the ability to enhance T cell proliferation when they are presented to CD70-expressing T cells either cross-linked with a secondary antibody or immobilized on tissue culture plates (Bowman *et al.*, 1994, *J Immunol* 152:1756-61; Brugnani, 1997, *Immunol Lett*
35 55:99-104). Such 'reverse signaling' has also been described in a subset of B chronic lymphocytic leukemia (B-CLL) cells, and CD70 can function as a receptor to transduce

5 signals to facilitate proliferation of PMA-stimulated purified B-CLL cells (Lens *et al.*, 1999, *supra*). These observations suggest situations in which engagement of CD27 and CD70 can result in the delivery of agonistic signals to both the CD27 and CD70 expressing cells.

[0010] The role of CD70/CD27 co-stimulation in cell-mediated autoimmune diseases has been investigated in a model of experimental autoimmune encephalomyelitis (EAE)
10 (Nakajima *et al.*, 2000, *J Neuroimmunol* 109:188-96). *In vivo* administration of a particular anti-mouse CD70 mAb (clone FR-70) markedly suppressed the onset of EAE by inhibiting antigen-induced TNF-alpha production without affecting T cell priming, Ig production or T_H1/T_H2 cell balance. However, such treatment had little efficacy in established disease. It has been reported that expression of CD70 on T cells was enhanced by TNF-alpha and IL-12
15 and down regulated by IL4 (Lens *et al.*, 1998, *supra*). Thus, the CD70/CD27 mediated T cell-T cell interactions may play a role in enhancing T_H1-mediated immune responses rather than T_H2-mediated responses. Supporting this hypothesis, the anti-mouse CD70 mAb FR-70 is also effective in inhibiting T_H1-mediated collagen-induced arthritis (Nakajima *et al.*, 2000, *supra*). In contrast, the same anti-mouse CD70 mAb did not show any efficacy in
20 modulating lupus in NZB/NZW F1 mice and experimental *Leishmania major* infection in susceptible BALB/c mice, both of which are predominantly T_H2-mediated autoimmune response (Nakajima *et al.*, 1997, *J Immunol* 158, 1466-72; Akiba *et al.*, 2000, *J Exp Med* 191:375-380).

[0011] The role of CD70 has not yet been investigated in acute graft versus host disease
25 (aGVHD), another T_H1-mediated immune response. GVHD is a major and often lethal consequence of allogeneic bone marrow transplantation (BMT) therapy that occurs when histocompatibility antigen differences between the BM donor and the recipient of the transplant are present (den Haan *et al.*, 1995, *Science* 268:1476). GVHD is caused by mature T cells present in the transplanted marrow, as well as other minor cell populations (Giralt and
30 Champlin, 1994, *Blood* 84:3603). It is noteworthy that CD70 has been detected *in vivo* on CD4+ cells in conditions characterized by allogeneic reaction, as in cases of maternal T cell engraftment in severe combined immune deficiency patients (Brugnoni *et al.*, *Immunol Lett* 55:99-104). Prophylaxis of GVHD is achieved by pan-T cell immunosuppressive agents such as cyclosporine, corticosteroids, or methotrexate. In addition to the lack of specificity,
35 these agents are also associated with significant adverse side effects. To limit these undesirable effects and the disruption of normal T cell functions, other therapeutic

5 interventions based on selective targeting of T cells directly participating in allo-recognition and graft rejection are much needed.

[0012] CD70 is a potentially useful target for antibody-directed immunotherapy. As indicated *supra*, CD70 has a restricted expression pattern in normal cells: CD70 expression is mostly restricted to recently antigen-activated T and B cells under physiological conditions, and its expression is down-regulated when antigenic stimulation ceases. The key role of CD70 is believed to be facilitating plasma cell differentiation and contributing to the generation and maintenance of long-term T cell memory. Further, evidence from animal models suggests that unregulated CD70/CD27 interaction may contribute to immunological disorders, and, in humans, experimental data have also pointed towards potential abnormal regulation of the CD70/CD27 pathway in T_H1-mediated immune disorders such as, *e.g.*, rheumatoid arthritis, psoriasis, and multiple sclerosis. It is of particular interest that CD70 is expressed on a variety of transformed cells including lymphoma B cells, Hodgkin and Reed-Sternberg cells, malignant cells of neural origin, and a number of carcinomas.

[0013] Several groups have demonstrated the inhibitory effect of anti-CD70 mAb in both *in vitro* models of lymphocyte activation and animal models of T_H1-mediated responses. The focus has been on the use of antibodies to block the CD70/CD27 co-stimulation pathway to achieve therapeutic efficacy. However, one main shortcoming of such an approach is the large number of signaling receptors, *e.g.*, the CD28/CD80/CD86 co-stimulatory pathway, known to participate in immunological diseases. Consequently, blocking one specific signaling pathway may only have minimal impact on disease development. This is supported by the observations that anti-CD70 mAb can only partially inhibit *in vitro* T cell activation induced by allogeneic stimulator cells (Hintzen *et al.*, 1995, *supra*) and an anti-CD70 mAb showed no therapeutic efficacy in EAE once the disease is established (Nakajima *et al.*, 2000, *supra*).

[0014] Thus, there is a need in the art for developing an approach for depleting or inhibiting the growth of CD70-expressing cells involved in cancers and/or immunological diseases by means other than or in addition to blocking the CD70/CD27 interaction. As CD70 is expressed on the surface of mature antigen presenting dendritic cells, activated T cells, and activated B cells, agents that can target and inhibit or deplete CD70⁺ cells may prove to be effective in the removal of antigen presenting cells presenting autoantigens and offending autoreactive activated T or B cells, as well as CD70-expressing tumor cells.

5 [0015] Approaches that have been used for increasing the therapeutic efficacy of antibodies are radiolabeling and combination with chemotherapy; however, these approaches include associated with undesirable side effects. For example, isotope therapy is associated with myelosuppression (Witzig, 2001, *Cancer Chemother Pharmacol* 48 (Suppl 1):S91-5), and
10 combining therapy with antibodies and chemotherapeutics is associated with immunosuppression. Further, isotopically labeled substances are difficult to produce, and patients often experience relapse after initial treatment with isotopically labeled substances.

[0016] Accordingly, there is a need for anti-CD70 ADCs that are constructed in such a manner so as to be capable exerting a clinically useful cytotoxic, cytostatic, or
15 immunosuppressive effect on CD70-expressing cells, particularly without exerting undesirable effects on non-CD70-expressing cells. Such compounds would be useful therapeutic agents against cancers that express CD70 or immune disorders that are mediated by CD70-expressing cells.

BRIEF SUMMARY OF THE INVENTION

20 [0017] The present invention provides antibody-drug conjugates (ADCs) and ADC derivatives and methods relating to the use of such conjugates to treat CD70-expressing cancers and immunological disorders. The antibody, or other targeting moiety in the ADC, binds to CD70. A drug conjugated to the antibody or targeting moiety exerts a cytotoxic, cytostatic, or immunosuppressive effect on CD70-expressing cells to treat or prevent
25 recurrence of CD70-expressing cancers or immunological disorders.

[0018] In one aspect, method are provided for the treatment of a CD70-expressing cancer in a subject. The methods generally include administering to the subject an effective amount of an antibody-drug conjugate. The antibody-drug conjugate includes an antibody that binds to CD70 ("CD70 antibody"). The antibody is conjugated to a drug that is a cytotoxic or
30 cytostatic agent. In certain embodiments, the antibody is the monoclonal antibody 1F6 or 2F2.

[0019] In other embodiments, the antibody binds to CD70 and competes for binding to CD70 with monoclonal antibody 1F6 or 2F2. Such a CD70 antibody can include at least one polypeptide region selected from (a) an H1 region having at least 80% sequence identity to
35 the amino acid sequence set forth in SEQ ID NO:6; (b) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:8; an H3 region having

5 at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:10; (d) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:16; (e) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:18; (f) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:20; (g) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:26; (h) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:28; (i) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:30; (j) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:36; (k) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:38; and (l) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:40.

[0020] In certain embodiments, the CD70 antibody includes the H1, H2, and H3 regions of (a), (b), and (c) (*supra*); or the H1, H2, and H3 regions of (g), (h), and (i) (*supra*). In additional embodiments, the H1, H2, H3, L1, L2 and L3 regions of the CD70 antibody includes polypeptide regions have the amino acid sequences set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10; SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20, respectively.

[0021] In additional embodiments, the H1, H2, and H3 regions correspond to (a), (b), and (c) (*supra*) and the L1, L2, and L3 regions correspond to (d), (e), and (f) (*supra*), respectively; or the H1, H2, and H3 regions correspond to (g), (h), and (i) (*supra*) and the L1, L2, and L3 regions correspond to (j), (k), and (l) (*supra*). In further embodiments, the CD70 antibody comprises a heavy chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO 22. The CD70 heavy chain variable region also can have the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:22.

[0022] In yet other embodiments, the heavy chain variable region has at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:22, and the light chain variable region has at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32. The light chain variable region also can have the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

[0023] In exemplary embodiments, the antibody-drug conjugate is 1F6-val-cit-AFP, 1F6-val-cit-MMAF, 1F6-val-cit-MMAE, 2F2-val-cit-AFP, 2F2-val-cit-MMAF, and 2F2-val-cit-MMAE.

5 [0024] The CD70 antibody can be, for example, a chimeric, humanized, or human antibody. In certain embodiments, the CD70 antibody is a chimeric antibody having a human constant region. The CD70 antibody can be monovalent or polyvalent.

[0025] Suitable cytotoxic agents can be, for example, an auristatin, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a
10 duocarmycin, a taxane, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid. In specific embodiments, the cytotoxic agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, paclitaxel, docetaxel, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, chaliceamicin, maytansine, DM-1, or netropsin. Other suitable cytotoxic agents include anti-tubulin agents,
15 such as an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin. In specific embodiments, the antitubulin agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcimid, estramustine, cemadotin, discodermolide, maytansine,
20 DM-1, or eleutherobin.

[0026] In antibody drug conjugates, the antibody can be conjugated directly to the cytotoxic agent or via a linker. Suitable linkers include, for example, cleavable and non-cleavable linkers. A cleavable linker is typically susceptible to cleavage under intracellular conditions. Suitable cleavable linkers include, for example, a peptide linker cleavable by an intracellular
25 protease, such as lysosomal protease or an endosomal protease. In exemplary embodiments, the linker can be a dipeptide linker, such as a valine-citrulline (val-cit) or a phenylalanine-lysine (phe-lys) linker. Other suitable linkers include linkers hydrolyzable at a pH of less than 5.5, such as a hydrazone linker. Additional suitable cleavable linkers include disulfide linkers.

30 [0027] An antibody-drug conjugate can be targeted to CD70-expressing cells, such as, for example, those of a kidney tumor, a B cell lymphoma, a thymic carcinoma, a nasopharyngeal carcinoma, or a brain tumor. The kidney tumor can be, for example, a renal cell carcinoma. The brain tumor can be, for example, a glioma, a glioblastoma, or a meningioma. The antibody-drug conjugates can be administered to humans or to non-human animals.

35 [0028] In another aspects, methods are provided for treating an immunological disorder. The methods generally include administering to a subject an amount effective of an antibody-

5 drug conjugate comprising an antibody that binds CD70. The antibody is conjugated to a cytotoxic agent or an immunosuppressive agent. In certain embodiments, the antibody is the monoclonal antibody 1F6 or 2F2.

[0029] In other embodiments, the antibody binds to CD70 and competes for binding to CD70 with monoclonal antibody 1F6 or 2F2. The CD70 antibody can include at least one
10 polypeptide region selected from (a) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:6; (b) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:8; an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:10; (d) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ
15 ID NO:16; (e) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:18; (f) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:20; (g) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:26; (h) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:28;
20 (i) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:30; (j) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:36; (k) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:38; and (l) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:40.

25 [0030] In certain embodiments, the CD70 antibody includes the H1, H2, and H3 regions of (a), (b), and (c) (*supra*); or the H1, H2, and H3 regions of (g), (h), and (i) (*supra*), respectively. In additional embodiments, the H1, H2, H3, L1, L2 and L3 regions of the CD70 antibody includes polypeptide regions have the amino acid sequences set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10; SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20,
30 respectively.

[0031] In additional embodiments, the H1, H2, and H3 regions correspond to (a), (b), and (c) (*supra*) and the L1, L2, and L3 regions correspond to (d), (e), and (f) (*supra*), respectively; or the H1, H2, and H3 regions correspond to (g), (h), and (i) (*supra*) and the L1, L2, and L3 regions correspond to (j), (k), and (l) (*supra*), respectively. In further
35 embodiments, the CD70 antibody comprises a heavy chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO

- 5 22. The CD70 heavy chain variable region also can have the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:22.

[0032] In yet other embodiments, the heavy chain variable region has at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:22, and the light chain variable region has at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:12 or SEQ
10 ID NO:32. The light chain variable region also can have the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

[0033] In exemplary embodiments, the antibody-drug conjugate is 1F6-val-cit-AFP, 1F6-val-cit-MMAF, 1F6-val-cit-MMAE, 2F2-val-cit-AFP, 2F2-val-cit-MMAF, and 2F2-val-cit-MMAE.

- 15 [0034] The CD70 antibody can be, for example, a chimeric, humanized, or human antibody. In certain embodiments, the CD70 antibody is a chimeric antibody having a human constant region. The CD70 antibody can be monovalent or polyvalent.

[0035] Suitable cytotoxic agents can be, for example, an auristatin, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a
20 duocarmycin, a taxane, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid. In specific embodiments, the drug is cytotoxic agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, paclitaxel, docetaxel, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, chaliceamicin, maytansine, DM-1, or netropsin. Other suitable cytotoxic agents include
25 anti-tubulin agents, such as an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin. In specific embodiments, the antitubulin agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcimid, estramustine, cemadotin,
30 discodermolide, maytansine, DM-1, or eleutherobin.

[0036] Suitable immunosuppressive agents include, for example, gancyclovir, etanercept, cyclosporine, tacrolimus, rapamycin, cyclophosphamide, azathioprine, mycophenolate mofetil, methotrexate, cortisol, aldosterone, dexamethasone, a cyclooxygenase inhibitor, a 5-lipoxygenase inhibitor, or a leukotriene receptor antagonist.

5 [0037] In antibody drug conjugates, the antibody can be conjugated directly to the cytotoxic agent or immunosuppressive agent, or via a linker. Suitable linkers include, for example, cleavable and non-cleavable linkers. A cleavable linker is typically susceptible to cleavage under intracellular conditions. Suitable cleavable linkers include, for example, a peptide linker cleavable by an intracellular protease, such as lysosomal protease or an
10 endosomal protease.

[0038] In exemplary embodiments the linker can be a dipeptide linker, such as a valine-citrulline (val-cit) or a phenylalanine-lysine (phe-lys) linker. Other suitable linkers include cleavable linker hydrolyzable at a pH of less than 5.5, such as a hydrazone linker. Further suitable cleavable linkers include disulfide linkers.

15 [0039] The ADC can be targeted to cells of an immunological disorder, such as, for example, Th1-mediated immunological disorders. Th1-mediated immunological disorders include, for example, rheumatoid arthritis, multiple sclerosis, psoriasis, Sjorgren's syndrome, Hashimoto's thyroiditis, Grave's disease, primary biliary cirrhosis, Wegener's granulomatosis, tuberculosis, or acute graft versus host disease. Other immunological
20 disorders include, for example, an activated B-lymphocyte disorder. The subject can be a human or a non-human animal.

[0040] In yet another aspect, antibody-drug conjugates are provided. The antibody-drug conjugates include an antibody that binds to CD70. The antibody is conjugated to a cytotoxic agent or an immunosuppressive agent. The antibody-drug conjugate can exert a cytotoxic or
25 cytostatic effect on a CD70-expressing cancer cell line, and/or a cytotoxic, cytostatic, or immunosuppressive effect on a CD70-expressing immune cell.

[0041] In certain embodiments, the antibody is the monoclonal antibody 1F6 or 2F2. In other embodiments, the antibody binds to CD70 and competes for binding to CD70 with monoclonal antibody 1F6 or 2F2. The CD70 antibody can include at least one polypeptide
30 region selected from (a) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:6; (b) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:8; an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:10; (d) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID
35 NO:16; (e) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:18; (f) an L3 region having at least 80% sequence identity to the

5 amino acid sequence set forth in SEQ ID NO:20; (g) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:26; (h) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:28; (i) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:30; (j) an L1 region having at least 80% sequence identity to the amino acid
10 sequence set forth in SEQ ID NO:36; (k) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:38; and (l) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:40.

[0042] In certain embodiments, the CD70 antibody includes the H1, H2, and H3 regions of (a), (b), and (c) (*supra*); or the H1, H2, and H3 regions of (g), (h), and (i) (*supra*),
15 respectively. In additional embodiments, the H1, H2, H3, L1, L2 and L3 regions of the CD70 antibody includes polypeptide regions have the amino acid sequences set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10; SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20, respectively.

[0043] In additional embodiments, the H1, H2, and H3 regions correspond to (a), (b), and
20 (c) (*supra*) and the L1, L2, and L3 regions correspond to (d), (e), and (f) (*supra*), respectively; or the H1, H2, and H3 regions correspond to (g), (h), and (i) (*supra*) and the L1, L2, and L3 regions correspond to (j), (k), and (l) (*supra*), respectively. In further embodiments, the CD70 antibody comprises a heavy chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO
25 22. The CD70 heavy chain variable region also can have the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:22.

[0044] In yet other embodiments, the heavy chain variable region has at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:22, and the light chain variable region has at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:12 or SEQ
30 ID NO:32. The light chain variable region also can have the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

[0045] In exemplary embodiments, the antibody-drug conjugate is 1F6-val-cit-AFP, 1F6-val-cit-MMAF, 1F6-val-cit-MMAE, 2F2-val-cit-AFP, 2F2-val-cit-MMAF, and 2F2-val-cit-MMAE.

5 [0046] The CD70 antibody can be, for example, a chimeric, humanized, or human antibody. In certain embodiments, the CD70 antibody is a chimeric antibody having a human constant region. The CD70 antibody can be monovalent or polyvalent.

[0047] Suitable cytotoxic agents can be, for example, an auristatin, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a
10 duocarmycin, a taxane, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid. In specific embodiments, the drug is cytotoxic agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, paclitaxel, docetaxel, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, chaliceamicin, maytansine, DM-1, or netropsin. Other suitable cytotoxic agents include
15 anti-tubulin agents, such as an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin. In specific embodiments, the antitubulin agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcemid, estramustine, cemadotin,
20 discodermolide, maytansine, DM-1, or eleutherobin.

[0048] Suitable immunosuppressive agents include, for example, gancyclovir, etanercept, cyclosporine, tacrolimus, rapamycin, cyclophosphamide, azathioprine, mycophenolate mofetil, methotrexate, cortisol, aldosterone, dexamethasone, a cyclooxygenase inhibitor, a 5-lipoxygenase inhibitor, or a leukotriene receptor antagonist.

25 [0049] In antibody drug conjugates, the antibody can be conjugated directly to the cytotoxic agent or immunosuppressive agent, or via a linker. Suitable linkers include, for example, cleavable and non-cleavable linkers. A cleavable linker is typically susceptible to cleavage under intracellular conditions. Suitable cleavable linkers include, for example, a peptide linker cleavable by an intracellular protease, such as lysosomal protease or an
30 endosomal protease.

[0050] Antibody drug conjugates can be targeted to CD70-expressing cancer cell, such as a renal cell carcinoma, a B cell lymphoma, or a glioblastoma. The antibody-drug conjugate also can be targeted to CD70-expressing immune cells, such as an activated T cell, an activated B cell, or a mature dendritic cell.

35 [0051] In yet additional aspects, pharmaceutical compositions are provided for the treatment of a CD70-expressing cancer or an immunological disorder. The pharmaceutical

- 5 compositions include an antibody-drug conjugate and at least one pharmaceutically compatible ingredient. In a related aspect, pharmaceutical kits are provided. The kits include a container containing an antibody-drug conjugate. The antibody-drug conjugate is typically lyophilized. The kits can further include a second container containing a pharmaceutically acceptable diluent.
- 10 [0052] The present invention may be more fully understood by reference to the following detailed description of the invention, non-limiting examples of specific embodiments of the invention and the appended figures.

BRIEF DESCRIPTION OF THE DRAWINGS

- 15 [0053] Figure 1: The 1F6 V_L and V_H cDNA and amino acid sequences. The coding and amino acid sequences for the light (V_L, upper 2 panels) and heavy chain (V_H, lower 2 panels) variable regions of 1F6 were determined. The complementarity determining regions (CDRs) for the V_L and V_H were identified according to criteria described in Kabat *et al.*, 1991, *Sequences of Proteins of Immunological Interest*, Washington, DC, US Department of Health and Public Services; Chothia and Lesk, 1987, *J Mol Biol* 196: 901-917. Amino acid residues corresponding to the CDRs are underlined. The signal peptide for the V_L and V_H are identified to be amino residues -20 to 0 and -19 to 0, respectively.

- [0054] Figure 2: The 2F2 V_L and V_H cDNA and amino acid sequences. The coding and amino acid sequences for the light (V_L, upper 2 panels) and heavy chain (V_H, lower 2 panels) variable regions of 2F2 were determined. The complementarity determining regions (CDRs) for the V_L and V_H were identified according to criteria described in Kabat *et al.*, 1991, *Sequences of Proteins of Immunological Interest*, Washington, DC, US Department of Health and Public Services; Chothia and Lesk, 1987, *J. Mol. Biol.* 196:901-917. Amino acid residues corresponding to the CDRs are underlined. The signal peptides for the V_L and V_H were identified to be amino residues -20 to 0 and -19 to 0, respectively.

[0055] Figure 3: Amino acid sequence comparison between the 1F6 and 2F2 CDRs. The amino acid sequences of 1F6 and 2F2 CDRs are aligned. Underlined residues represent conservative substitutions and boxed and italic residues represent divergent substitutions.

- [0056] Figure 4: Expression of CD70 on hematologic cell lines. CD70 expression on the surface of a panel of hematologic cell lines of different tissue origins and sources was

5 determined by flow cytometry. CD70 expression was expressed as binding ratios between the anti-CD70 mAb and control IgG.

[0057] Figure 5: Effects of anti-CD70 ADCs on the proliferation of transformed hematologic cell lines. A subset of the CD70⁺ cell lines shown in Figure 4 and the CD70⁻ Jurkat cell line were tested for their sensitivity toward 1F6 ADCs. Graded doses of the 1F6-vcMMAE, 1F6-vcAFP, or the corresponding non-binding control IgG (IgG) ADCs were
10 added to cells at the initiation of culture. Cells were exposed to the ADCs continuously for a total of 96 hours. Proliferation was determined by pulsing with ³H-TdR during the last 16 hours of incubation. The upper panels show the response of the WSU-NHL and Jurkat lines. The lower panel summarizes the responses by the tested cell lines to 1F6-vcMMAE and 1F6-vcAFP in form of IC₅₀s.
15

[0058] Figure 6: Cancer Profiling Array (CPA) analysis of the expression of CD70 message in multiple cancer types. Expression of CD70 in 241 tumor isolates and the corresponding adjacent normal tissues on a CPA was determined by hybridization with a CD70 cDNA probe. Thirteen tumor types are present on the array as indicated in the figure.
20 Each pair of dots represents a tumor sample (right side, T) and the corresponding normal tissue (left side, N). For some cDNA sets, as exemplified by the boxed sample set, the cDNA sample on the upper left spot is derived from the normal tissue, the upper right from the metastatic tumor, and the lower right from the primary tumor. Asterisks denote cDNA pairs in which CD70 transcript over-expression was detected in the tumor cDNA (tumor:normal
25 ratio of >2-fold, based on hybridization intensity).

[0059] Figure 7: Over-expression of CD70 transcripts in renal cell carcinoma (RCC) detected by quantitative PCR. CD70 expression in RCC cDNA samples was compared to that detected in a pool of 4 normal kidneys by quantitative PCR using an ABI PRISM 7000 Sequence Detection System and the $\Delta\Delta C_t$ method. Numbers represent the fold of CD70
30 transcript expression in RCC compared to that in normal kidneys.

[0060] Figures 8A and B: CD70 protein expression in frozen tumor sections of RCC. (A) Binding of the anti-CD70 mAb 2F2 (right column) and a control IgG (left column) to serial frozen tumor sections derived two RCC donors was determined by immunohistochemistry staining. Photomicrographs were taken under 40X magnification. (B) The same pair of
35 antibodies was used to stain serial frozen normal tissue sections adjacent to the tumor obtained from the same RCC donors used in (A).

5 [0061] Figures 9A and B: Expression of CD70 on renal cell carcinoma cell lines (RCC). CD70 expression on the surface of a panel of RCC was determined by flow cytometry. (A) The staining profiles of an anti-CD70 (open curves) and a control IgG (closed curves) on four representative RCC cell lines 786-O, Caki-1, Caki-2, and Hs 835.T and two normal renal tubule epithelial lines RPTEC and HRCE are shown. (B) CD70 expression on additional
10 RCC cell lines (CAL54, SK-RC-6, SK-RC-7, 786-O, A498, and ACHN) is compared to cell lines shown in (A). Mean fluorescence intensity of anti-CD70 staining and background fluorescence are plotted. Numbers on top of each bar indicate either mean fluorescence intensity obtained from anti-CD70 binding or background fluorescence.

[0062] Figure 10: Growth inhibitory and cytotoxicity activities of 1F6-vcAFP. 786-O or
15 Caki-1 cells were incubated with graded doses of 1F6-vcAFP or the non-binding control IgG-vcAFP. Cells were incubated for a total of 96 hours. ³H-thymidine incorporation (upper panel) and reduction of alamarBlue (lower panel) were used to determine proliferation and cell viability, respectively. Data points are expressed as percentages of the untreated control. Error bars represent the standard deviations of values obtained from quadruplicate wells.

20 [0063] Figure 11: Effects of anti-CD70 ADCs on the proliferation of RCC cell lines. Dose-response curves on growth inhibitory or cytotoxicity of 1F6 ADCs was determined as described in FIG 10 to obtain IC₅₀ values. The IC₅₀s of 1F6-vcMMAE, 1F6-vcMMAF, 1F6-vcAFP, or two control non-binding IgG (IgG) ADCs on the proliferation of CD70⁺ RCC lines (Caki-1, Caki-2, 786-O, 769-P, SK-RC-6, SK-RC-7, ACHN, CAL54, and A498), a CD70⁻
25 RCC line (Hs 835.T), and normal kidney tubule epithelial cell lines (RPTEC and HRCE) are summarized. Effects of 1F6-vcMMAF and 1F6-vcAFP on the viability of a subset of the cell lines are also summarized.

[0064] Figure 12: *In vivo* efficacy of 1F6-vcAFP in a RCC model. Tumors were initiated in nude mice by implanting tissue tumor blocks (30 mm³ in size) containing Caki-1 cells.
30 Treatment began when average tumor size within a treatment group was around 100 mm³. In the upper panel, treatment for the IgG and 1F6 groups began 17 days post implantation (filled arrow), while treatment for the 1F6-vcAFP and IgG-vcAFP groups began 25 days post implantation (open arrow). Indicated doses of either unconjugated mAbs or their ADCs were given for one course on a q4dx4 schedule. The lower panel shows results from a repeat
35 experiment when treatment began 14 days (filled arrow) after tumor implantation with 3 mg/kg and a q4dx4 schedule of 1F6-vcAFP or IgG-vcAFP.

5 [0065] Figure 13: Activation-induced CD70 expression on a T cell clone. A representative T cell clone C1A was activated by phytohemagglutinin A-L (PHA-L), irradiated feeder cells, IL-2, and IL-4. CD25 (left panel) and CD70 (right panel) expression on days 0, 2, 4, and 8 were determined by flow cytometric analysis.

[0066] Figure 14: Effects of anti-CD70 ADCs on the proliferation of activated T cell
10 clones. Resting T cell clones C2A, 40D8, and 4.01.1 were induced to express CD70 as described. At the peak of CD70 expression (2 days after activation), cells were harvested and treated with graded doses of 1F6-vcMMAE, 1F6-vcAFP, or the corresponding non-binding control IgG (IgG) ADCs as indicated in the figure in the presence of rhIL-2 and rhIL-4. Cells
15 were exposed to the ADCs continuously for an additional 72 hours. T cell proliferation was determined by pulsing with ^3H -TdR during the last 16 hours of incubation.

[0067] Figure 15: Effects of anti-CD70 ADCs on mixed lymphocyte reaction. An MLR was set up between PBMCs and irradiated allogeneic CESS cells. Expression of CD70 on activated allo-reactive T cells is shown on the upper panel. Binding of anti-CD70 mAb and control IgG is indicated by the open and closed curves, respectively. Graded doses of either
20 1F6-vcMMAE or 1F6-vcAFP were added to the culture at initiation. Cultures were exposed to the ADCs for a total of 120 hours. T cell proliferation was determined by pulsing with ^3H -thymidine during the last 16 hours of incubation (lower panel).

[0068] Figure 16: Effects of anti-CD70 or anti-CD70 ADC on antigen-induced T cell proliferation. T cells enriched for tetanus toxoid reactivity were stimulated with autologous
25 dendritic cells and tetanus toxoid in the presence of graded doses of 1F6, 1F6-vcMMAE, or 1F6-vcAFP for a total of 96 hours. T cell proliferation was determined by pulsing with ^3H -thymidine during the last 16 hours of incubation.

[0069] Figures 17A and B: CD70 induction during antigen-specific T cell expansion. PBMCs from a normal HLA-A0201 donor were stimulated with the M1 peptide derived from
30 the influenza virus matrix protein. (A) Expansion of M1 peptide-specific CD8^+ cells expressing TCR $\text{V}\beta 17$ chain was followed by flow cytometry (upper panels). The lower left panel shows the percentage of $\text{CD8}^+/\text{V}\beta 17^+$ cells expressing CD70 and the lower right panel shows the intensity of CD70 expression on the expanding $\text{CD8}^+/\text{V}\beta 17^+$ cells. (B) A representative example of specific CD70 induction on the expanding $\text{CD8}^+/\text{V}\beta 17^+$ after
35 stimulation with the M1 peptide for five days. Binding of the control IgG (open curves) and anti-CD70 mAb (closed curves) on the $\text{CD8}^+/\text{V}\beta 17^-$ or $\text{CD8}^+/\text{V}\beta 17^+$ cells are shown.

5 [0070] Figure 18: Specific deletion of antigen-specific CD8⁺/Vβ17⁺ T cells by anti-CD70 ADCs. PBMCs from a normal HLA-A0201 donor were stimulated with the M1 peptide for 5 days. 1F6 ADCs or control non-binding IgG (cIgG) ADCs were added to cells to a final concentration of 1 μg/ml. Total viable cell counts were conducted at 24, 48, and 76 hours post drug addition. The percentage of CD8⁺/Vβ17⁺ in each culture condition was determined
10 by flow cytometry as shown in Figure 17. Total numbers of CD8⁺/Vβ17⁺ for each culture condition are calculated (upper panel). The numbers of CD8⁺/Vβ17⁺ and CD8⁺/Vβ17⁻ cells in the 1F6-ADC treated cultures are expressed as percentages of the untreated control cultures (lower panel).

[0071] Figure 19: Dose-response comparison of anti-CD70 ADCs on deletion of antigen-specific CD8⁺/Vβ17⁺ cells. PBMCs from a normal HLA-A0201 donor were stimulated with the M1 peptide for 5 days. 1F6 ADCs or control non-binding IgG (cIgG) ADCs were given
15 to cells to the specified final concentrations. The percentages of CD8⁺/Vβ17⁺ cells 4 days post ADC addition was determined by flow cytometry as in Figure 17.

[0072] Figure 20: Limited effects of 1F6 ADCs on the proliferation capacity of the CD70⁻ bystander T cells. PBMCs from a normal HLA-A0201 donor were stimulated with the M1 peptide for 5 days. Cells were either allowed to expand or treated with 1 μg/ml 1F6-vcMMAF for an additional 4 days. Percentages of CD8⁺/Vβ17⁺ cells were determined by flow cytometry (left column). In order to assess the proliferation capacity of the CD70⁻ bystander, antigen non-specific T cells, Vβ17⁺ T cells were eliminated from the cultures by
20 immuno-depletion (center column). The resulting Vβ17⁻ cells were then stimulated with graded doses of immobilized anti-CD3 and anti-CD28 mAb for a total of 96 hours, and DNA synthesis was detected by pulsing with ³H-TdR during the last 18 hours of incubation.
25

DETAILED DESCRIPTION OF THE INVENTION

30 [0073] Provided are methods and compositions relating to the use of antibody-drug conjugates (ADCs) and ADC derivatives that bind to CD70. A drug conjugated to the antibody exerts a cytotoxic, cytostatic, or immunosuppressive effect on CD70-expressing cells to treat or immunological disorders or CD70-expressing cancers. In one aspect, the methods and compositions relate to antibodies and derivatives thereof that compete with
35 monoclonal antibody 1F6 or 2F2 for binding to CD70 and that, when conjugated to a cytotoxic agent, exert a cytotoxic or cytostatic effect to deplete or inhibit the proliferation of

- 5 CD70-expressing cells. Antibodies to CD70 which can be used in accordance with the methods and compositions described herein include monoclonal antibodies as well as chimeric, humanized, or human antibodies (*e.g.*, a humanized or chimeric form of 1F6 or 2F2), and such antibodies conjugated to cytotoxic or immunosuppressive agents such as, for example, chemotherapeutic drugs.
- 10 [0074] For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections which follow.

I. Definitions and Abbreviations

- [0075] Unless defined otherwise, all technical and scientific terms used herein have the same
15 meaning as commonly understood by one of ordinary skill in the art pertinent to the methods and compositions described. As used herein, the following terms and phrases have the meanings ascribed to them unless specified otherwise.

[0076] The term “inhibit” or “inhibition of” as used herein means to a reduce by a measurable amount, or to prevent entirely.

- 20 [0077] The term “agent” as used herein means an element, compound, or molecular entity, including, *e.g.*, a pharmaceutical, therapeutic, or pharmacologic compound. Agents can be natural or synthetic or a combination thereof. A “therapeutic agent” is an agent that exerts a therapeutic (*e.g.*, beneficial) effect on cancer cells or activated immune cells, either alone or in combination with another agent (*e.g.*, a prodrug converting enzyme in combination with a
25 prodrug). Typically, therapeutic agents useful in accordance with the methods and compositions described herein are those that exert a cytotoxic, cytostatic, or immunosuppressive effect.

- [0078] “Cytotoxic effect,” in reference to the effect of an agent on a cell, means killing of the cell. “Cytostatic effect” means an inhibition of cell proliferation. A “cytotoxic agent” means
30 an agent that has a cytotoxic or cytostatic effect on a cell, thereby depleting or inhibiting the growth of, respectively, cells within a cell population.

[0079] The term “deplete,” in the context of the effect of a CD70-targeting moiety-drug conjugate on CD70-expressing cells, refers to a reduction or elimination of the CD70-expressing cells.

5 [0080] The term “immunosuppressive agent” as used herein means an agent that inhibits the development or maintenance of an immunologic response. Such inhibition by an immunosuppressive agent can be effected by, for example, elimination of immune cells (*e.g.*, T or B lymphocytes); induction or generation of immune cells that can modulate (*e.g.*, down-regulate) the functional capacity of other cells; induction of an unresponsive state in immune
10 cells (*e.g.*, anergy); or increasing, decreasing or changing the activity or function of immune cells, including, for example, altering the pattern of proteins expressed by these cells (*e.g.*, altered production and/or secretion of certain classes of molecules such as cytokines, chemokines, growth factors, transcription factors, kinases, costimulatory molecules or other cell surface receptors, and the like). In typical embodiments, an immunosuppressive agent
15 has a cytotoxic or cytostatic effect on an immune cell that promotes an immune response.

[0081] “Immune cell” as used herein means any cell of hematopoietic lineage involved in regulating an immune response against an antigen (*e.g.*, an autoantigen). In typical embodiments, an immune cell is a T lymphocyte, a B lymphocyte, or a dendritic cell.

[0082] The term “polypeptide” refers to a polymer of amino acids and its equivalent and does
20 not refer to a specific length of a product; thus, “peptides” and “proteins” are included within the definition of a polypeptide. Also included within the definition of polypeptides are “antibodies” as defined herein. A “polypeptide region” refers to a segment of a polypeptide, which segment may contain, for example, one or more domains or motifs (*e.g.*, a polypeptide region of an antibody can contain, for example, one or more CDRs). The term “fragment”
25 refers to a portion of a polypeptide typically having at least 20 contiguous or at least 50 contiguous amino acids of the polypeptide. A “derivative” includes a polypeptide or fragment thereof having conservative amino acid substitutions relative to a second polypeptide; or a polypeptide or fragment thereof that is modified by covalent attachment of a second molecule such as, *e.g.*, by attachment of a heterologous polypeptide, or by
30 glycosylation, acetylation, phosphorylation, and the like. Further included within the definition of “polypeptide” is, for example, polypeptides containing one or more analogs of an amino acid (*e.g.*, unnatural amino acids and the like), polypeptides with unsubstituted linkages, as well as other modifications known in the art, both naturally and non-naturally occurring.

35 [0083] The term “antibody” as used herein refers to (a) immunoglobulin polypeptides and immunologically active portions of immunoglobulin polypeptides, *i.e.*, polypeptides of the

5 immunoglobulin family, or fragments thereof, that contain an antigen binding site that immunospecifically binds to a specific antigen (e.g., CD70), or (b) conservatively substituted derivatives of such immunoglobulin polypeptides or fragments that immunospecifically bind to the antigen (e.g., CD70). Antibodies are generally described in, for example, Harlow & Lane, *Antibodies: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1988).

10 [0084] In the context of immunoglobulin polypeptides or fragments thereof as defined above, “conservative substitution” means one or more amino acid substitutions that do not substantially reduce specific binding (e.g., as measured by the K_D) of the immunoglobulin polypeptide or fragment thereof to an antigen (i.e., substitutions that increase binding, that do not significantly alter binding, or that reduce binding by no more than about 40%, typically
15 no more than about 30%, more typically no more than about 20%, even more typically no more than about 10%, or most typically no more than about 5%, as determined by standard binding assays such as, e.g., ELISA).

[0085] An “antibody derivative” as used herein means an antibody, as defined above, that is modified by covalent attachment of a heterologous molecule such as, e.g., by attachment of a
20 heterologous polypeptide, or by glycosylation, acetylation or phosphorylation not normally associated with the antibody, and the like.

[0086] The term “monoclonal antibody” refers to an antibody that is derived from a single cell clone, including any eukaryotic or prokaryotic cell clone, or a phage clone, and not the method by which it is produced. Thus, the term “monoclonal antibody” as used herein is not
25 limited to antibodies produced through hybridoma technology.

[0087] The term “heterologous,” in the context of a polypeptide, means from a different source (e.g., a cell, tissue, organism, or species) as compared with another polypeptide, so that the two polypeptides are different. Typically, a heterologous polypeptide is from a different species.

30 [0088] As used herein, the term “functional,” in the context of an anti-CD70 antibody or derivative thereof to be used in accordance with the methods described herein, indicates that the antibody or derivative thereof is (1) capable of binding to CD70 and (2) depletes or inhibits the proliferation of CD70-expressing cells when conjugated to a cytotoxic agent, or has an immunosuppressive effect on an immune cell when conjugated to an
35 immunosuppressive agent.

5 [0089] The terms “identical” or “percent identity,” in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence. To determine the percent identity, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be
10 introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are
15 identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity = # of identical positions/total # of positions (*e.g.*, overlapping positions) x 100). In certain embodiments, the two sequences are the same length.

[0090] The term “substantially identical,” in the context of two nucleic acids or polypeptides,
20 refers to two or more sequences or subsequences that have at least 50%, at least 55%, at least 60%, or at least 65% identity; typically at least 70% or at least 75% identity; more typically at least 80% or at least 85% identity; and even more typically at least 90%, at least 95%, or at least 98% identity (as determined using one of the methods set forth *infra*).

[0091] “Similarity” or “percent similarity” in the context of two or more polypeptide
25 sequences, refer to two or more sequences or subsequences that have a specified percentage of amino acid residues that are the same or conservatively substituted when compared and aligned for maximum correspondence, as measured using one of the methods set forth *infra*. By way of example, a first amino acid sequence can be considered similar to a second amino acid sequence when the first amino acid sequence is at least 50%, 60%, 70%, 75%, 80%,
30 90%, or even 95% identical, or conservatively substituted, to the second amino acid sequence when compared to an equal number of amino acids as the number contained in the first sequence, or when compared to an alignment of polypeptides that has been aligned by a computer similarity program known in the art (*see infra*).

[0092] The terms “substantial similarity” or “substantial similarity,” in the context of
35 polypeptide sequences, indicates that a polypeptide region has a sequence with at least 70%, typically at least 80%, more typically at least 85%, and even more typically at least 90% or at

5 least 95% sequence similarity to a reference sequence. For example, a polypeptide is substantially similar to a second polypeptide, for example, where the two peptides differ by one or more conservative substitutions.

[0093] In the context of anti-CD70 antibodies or derivatives thereof, a protein that has one or more polypeptide regions substantially identical or substantially similar to one or more
10 antigen-binding regions (e.g., a heavy or light chain variable region, or a heavy or light chain CDR) of an anti-CD70 antibody retains specific binding to an epitope of CD70 recognized by the anti-CD70 antibody, as determined using any of various standard immunoassays known in the art or as referred to herein.

[0094] The determination of percent identity or percent similarity between two sequences can
15 be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul, 1993, *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, *et al.*, 1990, *J. Mol.*
20 *Biol.* 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to a nucleic acid encoding a protein of interest. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein of interest. To obtain gapped alignments for comparison purposes,
25 Gapped BLAST can be utilized as described in Altschul *et al.*, 1997, *Nucleic Acids Res.* 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Id.). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. (See, e.g., Internet web site address:
30 www.ncbi.nlm.nih.gov.) Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length
35 penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art and include ADVANCE and ADAM as described in Torellis and Robotti, 1994, *Comput. Appl. Biosci.* 10:3-5; and FASTA described in Pearson and

- 5 Lipman, 1988, *Proc. Natl. Acad. Sci.* 85:2444-8. Within FASTA, ktup is a control option that sets the sensitivity and speed of the search. If ktup=2, similar regions in the two sequences being compared are found by looking at pairs of aligned residues; if ktup=1, single aligned amino acids are examined. ktup can be set to 2 or 1 for protein sequences, or from 1 to 6 for DNA sequences. The default if ktup is not specified is 2 for proteins and 6 for DNA.
- 10 For a further description of FASTA parameters, *see, e.g.*, bioweb.pasteur.fr/docs/man/man/fasta.1.html#sect2, the contents of which are incorporated herein by reference.

[0095] Alternatively, protein sequence alignment may be carried out using the CLUSTAL W algorithm, as described by Higgins *et al.*, 1996, *Methods Enzymol.* 266:383-402.

- 15 [0096] As used herein, the terms "prevention" "prevent" refer to administration of an anti-CD70 antibody-drug conjugate (ADC) or ADC derivative to a subject before the onset of a clinical or diagnostic symptom of a CD70-expressing cancer or immunological disorder (*e.g.*, administration to an individual with a predisposition or at a high risk of acquiring the CD70-expressing cancer or immunological disorder) to (a) block the occurrence or onset of the
- 20 CD70-expressing cancer or immunological disorder, or one or more of clinical or diagnostic symptoms thereof, (b) inhibit the severity of onset of the CD70-expressing cancer or immunological disorder, or (c) to lessen the likelihood of the onset of the CD70-expressing cancer or immunological disorder.

- [0097] As used herein, the terms "treatment" or "treat" refer to slowing, stopping, or
- 25 reversing the progression of a CD70-expressing cancer or immunological disorder in a subject, as evidenced by a decrease or elimination of a clinical or diagnostic symptom of the disease, by administration of an anti-CD70 ADC or ADC derivative to the subject after the onset of the clinical or diagnostic symptom of the CD70-expressing cancer or immunological disorder at any clinical stage. Treatment can include, for example, a decrease in the severity
- 30 of a symptom, the number of symptoms, or frequency of relapse.

- [0098] The term "pharmaceutically acceptable" as used herein means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "pharmaceutically compatible ingredient" refers to a pharmaceutically
- 35 acceptable diluent, adjuvant, excipient, or vehicle with which a CD70-targeting moiety-drug conjugate is administered.

- 5 [0099] The term “effective amount,” in the context of the administration of an anti-CD70 ADC or ADC derivative, refers to the amount of the ADC or ADC derivative that is sufficient to inhibit the occurrence or ameliorate one or more clinical or diagnostic symptoms of a CD70-expressing cancer or immunological disorder in a subject. An effective amount of an agent is administered according to the methods described herein in an “effective regime.”
- 10 The term “effective regime” refers to a combination of amount of the agent and dosage frequency adequate to accomplish treatment or prevention of a CD70-expressing cancer or immunological disorder.

[0100] The abbreviation “AFP” refers to dimethylvaline-valine-dolaisoleuine- dolaproine-phenylalanine-p-phenylenediamine (*see* Formula XVI *infra*).

- 15 [0101] The abbreviation “MMAE” refers to monomethyl auristatin E (*see* Formula XI *infra*).

[0102] The abbreviation “AEB” refers to an ester produced by reacting auristatin E with paraacetyl benzoic acid (*see* Formula XX *infra*)

[0103] The abbreviation “AEVB” refers to an ester produced by reacting auristatin E with benzoylvaleric acid (*see* Formula XXI *infra*).

- 20 [0104] The abbreviation “MMAF” refers to dovaline-valine-dolaisoleunine-dolaproine-phenylalanine (*see* Formula IVIV *infra*).

[0105] The abbreviations “fk” and “phe-lys” refer to the linker phenylalanine-lysine.

[0106] The abbreviations “vc” and “val-cit” refer to the linker valine-citrulline.

25 II. Anti-CD70 Antibodies and Derivatives Thereof

- [0107] The methods and compositions described herein encompass the use of antibodies, or a derivatives thereof, as ADC’s that (a) specifically bind to CD70 and (b) when conjugated to a therapeutic agent, the therapeutic agent exerts a cytotoxic, cytostatic or immunosuppressive effect on CD70-expressing cancer cells or activated immune cells. As used herein, the term
- 30 “derivative,” in the context of an anti-CD70 antibody, refers to a molecule that (i) has an antigen-binding region of an anti-CD70 antibody, or a region derived therefrom (*e.g.*, by conservative substitution), and at least one polypeptide region or other moiety heterologous to the anti-CD70 antibody, and (ii) specifically binds to CD70 via the antigen-binding region or region derived therefrom. In specific embodiments, the anti-CD70 antibody is mAb 1F6

5 or 2F2 or a derivative thereof. In certain aspects, the anti-CD70 antibody or derivative thereof competes with monoclonal antibody 1F6 or 2F2 for binding to CD70.

[0108] In typical embodiments, the anti-CD70 antibody or derivative thereof, when conjugated to a cytotoxic agent, exerts a cytotoxic or cytostatic effect on CD70-expressing cancer cells, or, when conjugated to a cytotoxic or immunosuppressive agent, exerts a
10 cytotoxic, cytostatic, or immunosuppressive effect on activated lymphocytes or dendritic cells, for the treatment of a CD70-expressing cancer or an immunological disorder, respectively.

[0109] Anti-CD70 antibodies suitable for use in accordance with the present compositions and methods are typically monoclonal and can include, for example, chimeric (*e.g.*, having a
15 human constant region and mouse variable region), humanized, or human antibodies; single chain antibodies; or the like. The immunoglobulin molecules can be of any type (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY), class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule.

[0110] In certain embodiments, the antibody is an antigen-binding antibody fragment such
20 as, for example, a Fab, a F(ab'), a F(ab')₂, a Fd chain, a single-chain Fv (scFv), a single-chain antibody, a disulfide-linked Fv (sdFv), a fragment comprising either a V_L or V_H domain, or fragments produced by a Fab expression library, or a CD70-binding fragments of any of the above antibodies described *supra*. Antigen-binding antibody fragments, including single-chain antibodies, can comprise the variable region(s) alone or in combination with the
25 entirety or a portion of the following: hinge region, CH1, CH2, CH3 and CL domains. Also, antigen-binding fragments can comprise any combination of variable region(s) with a hinge region, CH1, CH2, CH3 and CL domains. Typically, the antibodies are human, rodent (*e.g.*, mouse and rat), donkey, sheep, rabbit, goat, guinea pig, camelid, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human
30 immunoglobulin and include antibodies isolated from human immunoglobulin libraries, from human B cells, or from animals transgenic for one or more human immunoglobulin, as described *infra* and, for example in U.S. Patent Nos. 5,939,598 and 6,111,166.

[0111] The antibodies may be monospecific, bispecific, trispecific, or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of CD70 or
35 may be specific for both CD70 as well as for a heterologous protein. (*See, e.g.*, PCT publications WO 93/17715; WO 92/08802; WO 91/00360; and WO 92/05793; Tutt *et al.*,

5 1991, *J Immunol* 147:60-69; U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; and 5,601,819; Kostelny *et al.*, 1992, *J Immunol* 148:1547-1553.) Multispecific antibodies, including bispecific and trispecific antibodies, useful for practicing the methods described herein are antibodies that immunospecifically bind to both CD70 (including but not limited to antibodies that have the CDRs and/or heavy chains of the monoclonal antibodies 2F2 and
10 1F6) and a second cell surface receptor or receptor complex, such as an immunoglobulin gene superfamily member, a TNF receptor superfamily member, an integrin, a cytokine receptor, a chemokine receptor, a major histocompatibility protein, a lectin (C-type, S-type, or I-type), or a complement control protein. In a typical embodiment, the binding of the portion of the multispecific antibody to the second cell surface molecule or receptor complex
15 enhances the cytotoxic or cytostatic effect of an anti-CD70 antibody-drug conjugate.

[0112] In certain specific embodiments, the anti-CD70 antibody is agonistic, non-agonistic or antagonistic with respect to CD70. In another specific embodiment, the anti-CD70 antibody does not block binding of CD70 ligand to CD70. In yet another embodiment, the anti-CD70 antibody or derivative thereof is a blocking antibody (*i.e.*, an antibody that blocks
20 the binding of CD27 to CD70).

[0113] In one aspect, an anti-CD70 antibody comprises one or more complementarity determining regions (CDRs) substantially identical or substantially similar to one or more CDR(s) of monoclonal antibody 1F6 (*see* Table 1). For example, the antibody can include a heavy chain CDR and/or a light chain CDR that is substantially identical or substantially
25 similar to a corresponding heavy chain CDR (H1, H2, or H3 regions) or corresponding light chain CDR (L1, L2, or L3 regions) of mAb 1F6 (SEQ ID NO:6; SEQ ID NO:8; SEQ ID NO:10; SEQ ID NO:16; SEQ ID NO:18; or SEQ ID NO:20, respectively). In typical embodiments, the anti-CD70 antibody has two or three heavy chain CDRs and/or two or three light chain CDRs that are substantially identical or substantially similar to
30 corresponding heavy and/or light chain CDRs of mAb 1F6. In specific embodiments, a CDR substantially identical or substantially similar to a heavy or light chain CDR of 1F6 has the amino acid sequence set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:16, SEQ ID NO:18, or SEQ ID NO:20.

[0114] For example, in certain embodiments, where an anti-CD70 antibody has at least one
35 heavy chain CDR substantially identical or substantially similar to a heavy chain CDR of

5 mAb 1F6, the antibody or derivative thereof further includes at least one light chain CDR that is substantially identical or substantially similar to a light chain CDR of mAb 1F6.

[0115] In certain typical embodiments, an anti-CD70 antibody includes a heavy or light chain variable domain, the variable domain having (a) a set of three CDRs substantially identical or substantially similar to corresponding CDRs of mAb 1F6, and (b) a set of four framework
10 regions. For example, an anti-CD70 antibody can include a heavy or light chain variable domain, the variable domain having (a) a set of three CDRs, in which the set of CDRs are from monoclonal antibody 1F6, and (b) a set of four framework regions, in which the set of framework regions are identical to or different from the set of framework regions in mAb 1F6.

15 [0116] In a specific embodiment, the anti-CD70 antibody includes a heavy chain variable region that is substantially identical or substantially similar to the heavy chain variable region of mAb 1F6 (*i.e.*, substantially identical or substantially similar to the amino acid sequences set forth in SEQ ID NO:2, *see* Table 1) and/or a light chain variable region that is substantially identical or substantially similar to the light chain variable regions of mAb 1F6
20 (*i.e.*, substantially identical or substantially similar to the amino acid sequences set forth in SEQ ID NO:12, *see* Table 1). For example, the antibody can include a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO:2 and, optionally, can further include a light chain variable region having the amino acid sequence set forth in SEQ ID NO:12. In one exemplary embodiment, the anti-CD70 antibody is mAb 1F6.

25 [0117] In another aspect, an anti-CD70 antibody comprises one or more CDRs substantially identical or substantially similar to one or more CDR(s) of monoclonal antibody 2F2 (*see* Table 1). For example, the antibody can include a heavy chain CDR and/or a light chain CDR that is substantially identical or substantially similar to a corresponding heavy chain CDR (H1, H2, or H3 regions) or corresponding light chain CDR (L1, L2, or L3 regions) of
30 mAb 2F2 (SEQ ID NO:26, SEQ ID NO:28; SEQ ID NO:30; SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:40). In typical embodiments, the anti-CD70 antibody has two or three heavy chain CDRs and/or two or three light chain CDRs that are substantially identical or substantially similar to corresponding heavy and/or light chain CDRs of mAb 2F2. In specific embodiments, a CDR substantially identical or substantially similar to a heavy or
35 light chain CDR of 2F2 has the amino acid sequence set forth in SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30; SEQ ID NO:36, SEQ ID NO:38, or SEQ ID NO:40.

5 [0118] For example, in certain embodiments, where an anti-CD70 antibody has at least one heavy chain CDR substantially identical or substantially similar to a heavy chain CDR of mAb 2F2, the antibody or derivative thereof further includes at least one light chain CDR that is substantially identical or substantially similar to a light chain CDR of mAb 2F2.

10 [0119] In certain typical embodiments, an anti-CD70 antibody includes a heavy or light chain variable domain, the variable domain having (a) a set of three CDRs substantially identical or substantially similar to corresponding CDRs of mAb 2F2, and (b) a set of four framework regions. For example, an anti-CD70 antibody can include a heavy or light chain variable domain, the variable domain having (a) a set of three CDRs, in which the set of CDRs are from monoclonal antibody 2F2, and (b) a set of four framework regions, in which the set of
15 framework regions are identical to or different from the set of framework regions in mAb 2F2.

[0120] In a specific embodiment, the anti-CD70 antibody includes a heavy chain variable region that is substantially identical or substantially similar to the heavy chain variable region of mAb 2F2 (*i.e.*, substantially identical or substantially similar to the amino acid sequences set forth in SEQ ID NO:22, *see* Table 1) and/or a light chain variable region that is
20 substantially identical or substantially similar to the light chain variable regions of mAb 2F2 (*i.e.*, substantially identical or substantially similar to the amino acid sequences set forth in SEQ ID NO:32, *see* Table 1). For example, the antibody can include a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO:22 and, optionally, can
25 further include a light chain variable region having the amino acid sequence set forth in SEQ ID NO:32. In one exemplary embodiment, the anti-CD70 antibody is mAb 2F2.

[0121] A table indicating the region of 1F6 or 2F2 to which each SEQ ID NO. corresponds is provided below:

Table 1

MOLECULE	NUCLEOTIDE OR AMINO ACID	SEQ ID NO
1F6 Heavy Chain Variable Region	Nucleotide	1
1F6 Heavy Chain Variable Region	Amino Acid	2
1F6 Heavy Chain Signal Peptide	Nucleotide	3
1F6 Heavy Chain Signal Peptide	Amino Acid	4
1F6 Heavy Chain-CDR1(H1)	Nucleotide	5

Table 1

MOLECULE	NUCLEOTIDE OR AMINO ACID	SEQ ID NO
1F6 Heavy Chain-CDR1(H1)	Amino Acid	6
1F6 Heavy Chain-CDR2(H2)	Nucleotide	7
1F6 Heavy Chain-CDR2(H2)	Amino Acid	8
1F6 Heavy Chain-CDR3(H3)	Nucleotide	9
1F6 Heavy Chain-CDR3(H3)	Amino Acid	10
1F6 Light Chain Variable Region	Nucleotide	11
1F6 Light Chain Variable Region	Amino Acid	12
1F6 Light Chain Signal Peptide	Nucleotide	13
1F6 Light Chain Signal Peptide	Amino Acid	14
1F6 Light Chain-CDR1(L1)	Nucleotide	15
1F6 Light Chain-CDR1(L1)	Amino Acid	16
1F6 Light Chain-CDR2(L2)	Nucleotide	17
1F6 Light Chain-CDR2(L2)	Amino Acid	18
1F6 Light Chain-CDR3(L3)	Nucleotide	19
1F6 Light Chain-CDR3(L3)	Amino Acid	20
2F2 Heavy Chain Variable Region	Nucleotide	21
2F2 Heavy Chain Variable Region	Amino Acid	22
2F2 Heavy Chain Signal Peptide	Nucleotide	23
2F2 Heavy Chain Signal Peptide	Amino Acid	24
2F2 Heavy Chain-CDR1(H1)	Nucleotide	25
2F2 Heavy Chain-CDR1(H1)	Amino Acid	26
2F2 Heavy Chain-CDR2(H2)	Nucleotide	27
2F2 Heavy Chain-CDR2(H2)	Amino Acid	28
2F2 Heavy Chain-CDR3(H3)	Nucleotide	29
2F2 Heavy Chain-CDR3(H3)	Amino Acid	30
2F2 Light Chain Variable Region	Nucleotide	31
2F2 Light Chain Variable Region	Amino Acid	32
2F2 Light Chain Signal Peptide	Nucleotide	33
2F2 Light Chain Signal Peptide	Amino Acid	34
2F2 Light Chain-CDR1(L1)	Nucleotide	35
2F2 Light Chain-CDR1(L1)	Amino Acid	36
2F2 Light Chain-CDR2(L2)	Nucleotide	37

Table 1

MOLECULE	NUCLEOTIDE OR AMINO ACID	SEQ ID NO
2F2 Light Chain-CDR2(L2)	Amino Acid	38
2F2 Light Chain-CDR3(L3)	Nucleotide	39
2F2 Light Chain-CDR3(L3)	Amino Acid	40

- 5 [0122] Anti-CD70 antibodies useful in the methods set forth herein may also be described or specified in terms of their binding affinity to CD70. Typical binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

- [0123] The antibodies that may be used in the treatment of immunological disorders or CD70-expressing cancers can be generated by any suitable method known in the art. For example, monoclonal antibodies can be prepared using a wide variety of techniques including, *e.g.*, the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. Hybridoma techniques are generally discussed in, for example, Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed., 1988); and Hammerling, *et al.*, In *Monoclonal Antibodies and T-Cell Hybridomas*, pp. 563-681 (Elsevier, N.Y., 1981). Examples of phage display methods that can be used to make the anti-CD70 antibodies include, *e.g.*, those disclosed in Brinkman *et al.*, 1995, *J Immunol Methods* 182:41-50; Ames *et al.*, 1995, *J Immunol Methods* 184:177-186; Kettleborough *et al.*, 1994, *Eur J Immunol* 24:952-958; Persic *et al.*, 1997, *Gene* 187:9-18; Burton *et al.*, 1994, *Advances in Immunology* 57:191-280; PCT Application No. PCT/GB91/01 134; PCT Publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108 (the disclosures of which are incorporated by reference herein).

- [0124] Techniques for generating antibody fragments that recognize specific epitopes are also generally known in the art. For example, Fab and $F(ab')_2$ fragments can be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce $F(ab')_2$ fragments). $F(ab')_2$ fragments contain

- 5 the variable region, the light chain constant region and the CH 1 domain of the heavy chain. Techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using, e.g., methods disclosed in PCT publication WO 92/22324; Mullinax *et al.*, 1992, *BioTechniques* 12(6):864-869; and Sawai *et al.*, 1995, *AJRI* 34:26-34; and Better *et al.*, 1988, *Science* 240:1041-1043 (the disclosures of which are incorporated by reference herein).
- 10 [0125] Examples of techniques that can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston *et al.*, 1991, *Methods in Enzymology* 203:46-88; Shu *et al.*, 1993, *Proc Natl Acad Sci USA* 90:7995-7999; and Skerra *et al.*, 1988, *Science* 240:1038-1040.
- 15 [0126] In certain embodiments, the anti-CD70 antibody is a chimeric antibody. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as for example antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. (See e.g., Morrison, *Science*, 1985, 229:1202; Oi *et al.*, 1986, *BioTechniques* 4:214; Gillies *et al.*, 1989, *J. Immunol. Methods* 20 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397.)
- [0127] An anti-CD70 antibody can also be a humanized antibody. Humanized antibodies are antibody molecules that bind the desired antigen and have one or more CDRs from a non-human species, and framework and constant regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted 25 with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen *et al.*, U.S. Patent No. 30 5,585,089; Riechmann *et al.*, 1988, *Nature* 332:323). Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 0 239 400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 0 592 106; EP 0 519 596; Padlan, *Molecular Immunology*, 1991, 28(4/5):489-498; Studnicka *et al.*, 1994, *Protein Engineering* 7(6):805-814; Roguska *et al.*, 1994, *PNAS* 91:969-973), and chain shuffling (U.S. Patent No. 5,565,332) (all of these 35 references are incorporated by reference herein).

5 [0128] In yet other embodiments, the anti-CD70 antibody is a human antibody. Human antibodies can be made by a variety of methods known in the art including, *e.g.*, phage display methods (*see supra*) using antibody libraries derived from human immunoglobulin sequences. *See also, e.g.*, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735,
10 and WO 91/10741. In addition, a human antibody recognizing a selected epitope can be generated using a technique referred to as "guided selection," in which a selected non-human monoclonal antibody, *e.g.*, a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope (*see, e.g.*, Jespers *et al.*, 1994, *Bio/technology* 12:899-903). Human antibodies can also be produced using transgenic mice that express
15 human immunoglobulin genes. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. For an overview of this technology for producing human antibodies, *see* Lonberg and Huszar, 1995, *Int. Rev. Immunol.* 13:65-93. For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing
20 such antibodies, *see, e.g.*, PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598, 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598. In addition, companies such as Abgenix, Inc. (Fremont, CA) and Medarex (Princeton, NJ) can be engaged to provide human antibodies directed against a selected
25 antigen using technology similar to that described above.

[0129] As set forth *supra*, a derivative of an anti-CD70 antibody can also be used in the practice of present methods. Generally, an anti-CD70 antibody derivative comprises an anti-CD70 antibody (including, *e.g.*, an antigen-binding fragment or conservatively substituted polypeptides) and at least one polypeptide region or other moiety heterologous to the anti-
30 CD70 antibody. For example, an anti-CD70 antibody can be modified, *e.g.*, by the covalent attachment of any type of molecule, such that covalent attachment does not prevent the antibody derivative from specifically binding to CD70 via the antigen-binding region or region derived therefrom, or the conjugated drug from exerting (a) a cytostatic or cytotoxic effect on CD70-expressing cancer cells, or (b) a cytostatic, cytotoxic, or immunosuppressive
35 effect on activated lymphocytes. Typical modifications include, *e.g.*, glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein,

5 and the like. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, *etc.* Additionally, the derivative may contain one or more non-classical amino acids.

[0130] In certain embodiments, the antibody derivative is a multimer, such as, for example, a
10 dimer, comprising one or more monomers, where each monomer includes (i) an antigen-binding region of an anti-CD70 antibody, or a polypeptide region derived therefrom (such as, *e.g.*, by conservative substitution of one or more amino acids), and (ii) a multimerizing (*e.g.*, dimerizing) polypeptide region, such that the antibody derivative forms multimers (*e.g.*, homodimers) that specifically bind to CD70. In typical embodiments, an antigen binding
15 region of an anti-CD70 antibody, or a polypeptide region derived therefrom, is recombinantly or chemically fused with a heterologous protein, wherein the heterologous protein comprises a dimerization or multimerization domain. Prior to administration of the antibody derivative to a subject for the purpose of treating or preventing immunoglocal disorders or CD70-expressing cancers, the derivative is subjected to conditions that allows formation of a
20 homodimer or heterodimer. A heterodimer, as used herein, may comprise identical dimerization domains but different CD70 antigen-binding regions, identical CD70 antigen-binding regions but different dimerization domains, or different CD70 antigen-binding regions and dimerization domains.

[0131] Typical dimerization domains are those that originate from transcription factors. In
25 one embodiment, the dimerization domain is that of a basic region leucine zipper ("bZIP"). (See C. Vinson *et al.*, 1989, *Science* 246:911-916). Useful leucine zipper domains include, for example, those of the yeast transcription factor GCN4, the mammalian transcription factor CCAAT/enhancer-binding protein C/EBP, and the nuclear transform in oncogene products, Fos and Jun. (See Landschultz *et al.*, 1988, *Science* 240:1759-1764; Baxeavanis and Vinson,
30 1993, *Curr. Op. Gen. Devel.* 3:278-285; O'Shea *et al.*, 1989, *Science* 243:538-542.) In another embodiment, the dimerization domain is that of a basic-region helix-loop-helix ("bHLH") protein. (See Murre *et al.*, 1989, *Cell* 56:777-783. See also Davis *et al.*, 1990, *Cell* 60:733-746; Voronova and Baltimore, 1990, *Proc Natl Acad Sci USA* 87:4722-4726.) Particularly useful hHLH proteins are myc, max, and mac.

35 [0132] In yet other embodiments, the dimerization domain is an immunoglobulin constant region such as, for example, a heavy chain constant region or a domain thereof (*e.g.*, a CH1

- 5 domain, a CH2 domain, or a CH3 domain). (See, e.g., U.S. Patent Nos. 5,155,027; 5,336,603; 5,359,046; and 5,349,053; EP 0 367 166; WO 96/04388.)

[0133] Heterodimers are known to form between Fos and Jun (Bohmann *et al.*, 1987, *Science* 238:1386-1392), among members of the ATF/CREB family (Hai *et al.*, 1989, *Genes Dev.* 3:2083-2090), among members of the C/EBP family (Cao *et al.*, 1991, *Genes Dev.* 5:1538-1552; Williams *et al.*, 1991, *Genes Dev.* 5:1553-1567; Roman *et al.*, 1990, *Genes Dev.* 4:1404-1415), and between members of the ATF/CREB and Fos/Jun families Hai and Curran, 1991, *Proc Natl Acad Sci USA* 88:3720-3724). Therefore, when a CD70-binding protein-drug conjugate is administered to a subject as a heterodimer comprising different dimerization domains, any combination of the foregoing may be used.

15 [0134] In other embodiments, an anti-CD70 antibody derivative is an anti-CD70 antibody conjugated to a second antibody (an "antibody heteroconjugate") (see U.S. Patent No. 4,676,980). Heteroconjugates useful for practicing the present methods comprise an antibody that binds to CD70 (e.g., an antibody that has the CDRs and/or heavy chains of the monoclonal antibodies 2F2 or 1F6) and an antibody that binds to a surface receptor or
20 receptor complex, such as an immunoglobulin gene superfamily member, a TNF receptor superfamily member, an integrin, a cytokine receptor, a chemokine receptor, a major histocompatibility protein, a lectin (C-type, S-type, or I-type), or a complement control protein.

[0135] In certain embodiments, the anti-CD70 antibody or derivative thereof competitively
25 inhibits binding of mAb 1F6 or 2F2 to CD70, as determined by any method known in the art for determining competitive binding (such as, e.g., the immunoassays described herein). In typical embodiments, the antibody competitively inhibits binding of 1F6 or 2F2 to CD70 by at least 50%, more typically at least 60%, yet more typically at least 70%, and most typically at least 75%. In other embodiments, the antibody competitively inhibits binding of 1F6 or
30 2F2 to CD70 by at least 80%, at least 85%, at least 90%, or at least 95%.

[0136] Antibodies can be assayed for specific binding to CD70 by any of various known methods. Immunoassays which can be used include, for example, competitive and non-competitive assay systems using techniques such as Western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays,
35 immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays,

- 5 immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well-known in the art. (*See, e.g., Ausubel et al., eds., Short Protocols in Molecular Biology* (John Wiley & Sons, Inc., New York, 4th ed. 1999); Harlow & Lane, *Using Antibodies: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1999).)
- 10 [0137] Further, the binding affinity of an antibody to CD70 and the off-rate of an antibody CD70 interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled CD70 (*e.g., ^3H or ^{125}I*) with the antibody of interest in the presence of increasing amounts of unlabeled CD70, and the detection of the antibody bound to the labeled CD70. The affinity
- 15 of the antibody for CD70 and the binding off-rates can then be determined from the data by Scatchard plot analysis. Competition with a second antibody (such as, *e.g.,* mAb 1F6 or 2F2) can also be determined using radioimmunoassays. In this case, CD70 is incubated with the antibody of interest conjugated to a labeled compound (*e.g., ^3H or ^{125}I*) in the presence of increasing amounts of an unlabeled second antibody. Alternatively, the binding affinity of an
- 20 antibody to CD70 and the on- and off-rates of an antibody-CD70 interaction can be determined by surface plasmon resonance.

- [0138] In accordance with the methods described herein, anti-CD70 antibodies or derivatives thereof, when conjugated to a therapeutic agent, can be internalized and accumulate within a CD70-expressing cell, where the therapeutic agent exerts an effect (*e.g.,* a cytotoxic,
- 25 cytostatic, or immunosuppressive effect). In additional embodiments, anti-CD70 antibodies or derivatives thereof, when conjugated to a therapeutic agent, can be targeted to and accumulate on the membrane of a CD70-expressing cell, where the therapeutic agent exerts an effect (*e.g.,* a cytotoxic, cytostatic, or immunosuppressive effect). In yet other embodiments, anti-CD70 antibodies or derivatives thereof, when conjugated to a therapeutic
- 30 agent, can be targeted to a biological molecules in a cell (*e.g.,* an inflammatory agent) and accumulate at or adjacent cells secreting or binding the biological molecule, where the therapeutic agent exerts an effect (*e.g.,* a cytotoxic, cytostatic, or immunosuppressive effect).

- [0139] Whether a given anti-CD70 antibody or derivative, when conjugated to a therapeutic agent, exerts a corresponding therapeutic effect upon binding a CD70-expressing
- 35 cell can be readily determined, *e.g.,* by (1) incubating CD70-expressing cells independently with the anti-CD70 antibody or derivative thereof, (2) incubating the cells with a secondary

5 reagent that is conjugated to the therapeutic agent and that specifically binds to the antibody or derivative thereof, and (3) assaying the cells for the corresponding therapeutic effect. Multiple antibodies or antibody derivatives can be readily evaluated via such assays using a secondary reagent that specifically binds a polypeptide region shared by each antibody or derivative thereof (e.g., an anti-Ig antibody). For example, an anti-CD70 mAb that binds
10 CD70 and exerts a cytotoxic effect when conjugated to a cytotoxic agent (e.g., an auristatin such as, for example, AFP, MMAF, or MMAE) can be identified by an indirect immunotoxicity assay such as, for example, described by Chun *et al.*, 2003, *Supplement to Clinical Cancer Research*, Vol. 9. Briefly, the cytotoxic agent is conjugated to a secondary antibody (e.g., for murine mAbs, a polyclonal anti-mouse IgG); CD70-expressing cells are
15 incubated with both the primary and cytotoxic agent-conjugated secondary antibody (e.g., in 96-well plates, using hybridoma supernatant for the primary antibody); and primary antibody-dependent cytotoxicity is assessed in a standard cytotoxicity assay (e.g., an MTT cell viability assay). (*See id.*)

[0140] The anti-CD70 antibodies and derivatives thereof that are useful in the present
20 methods can be produced by any method known in the art for the synthesis of proteins, typically, e.g., by recombinant expression techniques. Recombinant expression of an antibody or derivative thereof that binds to CD70 and depletes or inhibits the proliferation of CD70-expressing cells requires construction of an expression vector containing a nucleic acid that encodes the antibody or derivative thereof. Once a nucleic acid encoding such a protein
25 has been obtained, the vector for the production of the protein molecule may be produced by recombinant DNA technology using techniques well known in the art. Standard techniques such as, for example, those described in Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 3rd ed., 2001); Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor
30 Laboratory Press, Cold Spring Harbor, N.Y., 2nd ed., 1989); *Short Protocols in Molecular Biology* (Ausubel *et al.*, John Wiley & Sons, New York, 4th ed., 1999); and Glick & Pasternak, *Molecular Biotechnology: Principles and Applications of Recombinant DNA* (ASM Press, Washington, D.C., 2nd ed., 1998) can be used for recombinant nucleic acid methods, nucleic acid synthesis, cell culture, transgene incorporation, and recombinant
35 protein expression.

[0141] For example, for recombinant expression of an anti-CD70 antibody, an expression vector may encode a heavy or light chain thereof, or a heavy or light chain variable domain,

5 operably linked to a promoter. An expression vector may include, for example, the nucleotide sequence encoding the constant region of the antibody molecule (*see, e.g.*, PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464), and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain. The expression vector is transferred to a host cell by
10 conventional techniques, and the transfected cells are then cultured by conventional techniques to produce the anti-CD70 antibody. In typical embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains can be co-expressed in the host cell for expression of the entire immunoglobulin molecule.

[0142] A variety of prokaryotic and eukaryotic host-expression vector systems can be
15 utilized to express an anti-CD70 antibody or derivative thereof. Typically, eukaryotic cells, particularly for whole recombinant anti-CD70 antibody molecules, are used for the expression of the recombinant protein. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus, is an effective expression system for
20 the production of anti-CD70 antibodies and derivatives thereof (*see, e.g.*, Foecking *et al.*, 1986, *Gene* 45:101; Cockett *et al.*, 1990, *Bio/Technology* 8:2).

[0143] Other host-expression systems include, for example, plasmid-based expression systems in bacterial cells (*see, e.g.*, Ruther *et al.*, 1983, *EMBO* 1,2:1791; Inouye & Inouye, 1985, *Nucleic Acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.*
25 24:5503-5509); insect systems such as, *e.g.*, the use of *Autographa californica* nuclear polyhedrosis virus (AcNPV) expression vector in *Spodoptera frugiperda* cells; and viral-based expression systems in mammalian cells, such as, *e.g.*, adenoviral-based systems (*see, e.g.*, Logan & Shenk, 1984, *Proc. Natl. Acad. Sci. USA* 81:355-359; Bittner *et al.*, 1987, *Methods in Enzymol.* 153:51-544).

30 [0144] In addition, a host cell strain can be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing (*e.g.*, glycosylation, phosphorylation, and cleavage) of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular
35 machinery for proper processing of the primary transcript and gene product can be used.

- 5 Such mammalian host cells include, for example, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, and W138.

[0145] A stable expression system is typically used for long-term, high-yield production of recombinant anti-CD70 antibody or derivative thereof. For example, cell lines that stably express the anti-CD70 antibody or derivative thereof can be engineered by transformation of
10 host cells with DNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites) and a selectable marker, followed by growth of the transformed cells in a selective media. The selectable marker confers resistance to the selection and allows cells to stably integrate the DNA into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell
15 lines. A number of selection systems can be used, including, for example, the herpes simplex virus thymidine kinase, hypoxanthineguanine phosphoribosyltransferase, and adenine phosphoribosyltransferase genes, which can be employed in *tk*⁻, *hgp*⁺ or *ap*⁺ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: *dhfr*, which confers resistance to methotrexate; *gpt*, which confers resistance
20 to mycophenolic acid; *neo*, which confers resistance to the aminoglycoside G-418; and *hygro*, which confers resistance to hygromycin. Methods commonly known in the art of recombinant DNA technology can be routinely applied to select the desired recombinant clone, and such methods are described, for example, in *Current Protocols in Molecular Biology* (Ausubel *et al.* eds., John Wiley & Sons, N.Y., 1993); Kriegler, *Gene Transfer and*
25 *Expression, A Laboratory Manual* (Stockton Press, N.Y., 1990); *Current Protocols in Human Genetics* (Dracopoli *et al.* eds., John Wiley & Sons, N.Y., 1994, Chapters 12 and 13); and Colberre-Garapin *et al.*, 1981, *J. Mol. Biol.* 150:1.

[0146] The expression levels of an antibody or derivative can be increased by vector amplification. (See generally, *e.g.*, Bebbington & Hentschel, *The Use of Vectors Based on*
30 *Gene Amplification for the Expression of Cloned Genes in Mammalian Cells in DNA Cloning, Vol.3* (Academic Press, New York, 1987).) When a marker in the vector system expressing an anti-CD70 antibody or derivative thereof is amplifiable, an increase in the level of inhibitor present in host cell culture media will select host cells that have increased copy number of a marker gene conferring resistance to the inhibitor. The copy number of an
35 associated antibody gene will also be increased, thereby increasing expression of the antibody or derivative thereof (see Crouse *et al.*, 1983, *Mol. Cell. Biol.* 3:257).

- 5 [0147] Where the anti-CD70 antibody comprises both a heavy and a light chain or derivatives thereof, the host cell may be co-transfected with two expression vectors, the first vector encoding the heavy chain protein and the second vector encoding the light chain protein. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain proteins. Alternatively, a single vector may be used
- 10 which encodes, and is capable of expressing, both heavy and light chain proteins. In such situations, the light chain is typically placed before the heavy chain to avoid an excess of toxic free heavy chain (*see* Proudfoot, 1986, *Nature* 322:52; Kohler, 1980, *Proc. Natl. Acad. Sci. USA* 77:2197). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.
- 15 [0148] Once an anti-CD70 antibody or derivative thereof has been produced (*e.g.*, by an animal, chemical synthesis, or recombinant expression), it can be purified by any suitable method for purification of proteins, including, for example, by chromatography (*e.g.*, ion exchange or affinity chromatography (such as, for example, Protein A chromatography for purification of antibodies having an intact Fc region)), centrifugation, differential solubility,
- 20 or by any other standard technique for the purification of proteins. An anti-CD70 antibody or derivative thereof can, for example, be fused to a marker sequence, such as a peptide, to facilitate purification by affinity chromatography. Suitable marker amino acid sequences include, *e.g.*, a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), and the "HA" tag, which corresponds to an
- 25 epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, 1984, *Cell* 37:767), and the "flag" tag.
- [0149] Once an anti-CD70 antibody or derivative thereof is produced, its ability to exert a cytostatic or cytotoxic effect on CD70-expressing cancer cells (when conjugated to a cytotoxic agent) or an immunosuppressive effect on a CD70-expressing immune cell (when
- 30 conjugated to an immunosuppressive agent) is determined by the methods described *infra* or as known in the art.

III. Anti-CD70 Antibody-Drug Conjugates

- [0150] Compositions useful in the treatment of a CD70-expressing cancer or an
- 35 immunological disorder comprise anti-CD70 antibody-drug conjugates (ADCs) or anti-CD70 ADC derivatives. An "anti-CD70 ADC" as used herein refers to an anti-CD70 antibody (as

5 described in Section II, *supra*) conjugated to a therapeutic agent. An “anti-CD70 derivative” as used herein refers to derivative of an anti-CD70 antibody (as describein in Section II, *supra*) conjugated to a therapeutic agent. In certain embodiments, the ADC comprises an anti-CD70 antibody (*e.g.*, mAb 1F6 or 2F2 or a fragment or derivative thereof, including, for example, a chimeric or humanized form thereof). The ADCs or ADC derivatives as
10 described herein produce clinically beneficial effects on CD70-expressing cells when administered to a subject with a CD70-expressing cancer or an immunological disorder, typically when administered alone but also in combination with other therapeutic agents.

[0151] In typical embodiments, the anti-CD70 antibody or derivative thereof is conjugated to a cytotoxic or immunosuppressive agent, such that the resulting ADC or ADC derivative
15 exerts a cytotoxic or cytostatic effect on a CD70-expressing cancer cell, or a cytotoxic, cytostatic, or immunosuppressive effect on an immune cell (*e.g.*, an activated lymphocyte or dendritic cell) when taken up or internalized by the cell. Particularly suitable moieties for conjugation to antibodies or antibody derivatives are chemotherapeutic agents, prodrug converting enzymes, radioactive isotopes or compounds, or toxins. For example, an anti-
20 CD70 antibody or derivative thereof can be conjugated to a cytotoxic agent such as a chemotherapeutic agent (*see infra*), or a toxin (*e.g.*, a cytostatic or cytocidal agent such as, *e.g.*, abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin). Examples of additional agents that are useful for conjugating to the anti-CD70 molecules are provided *infra*.

[0152] In other embodiments, the anti-CD70 antibody or derivative thereof is conjugated to a
25 pro-drug converting enzyme. The pro-drug converting enzyme can be recombinantly fused to the antibody or derivative thereof or chemically conjugated thereto using known methods. Exemplary pro-drug converting enzymes are carboxypeptidase G2, beta-glucuronidase, penicillin-V-amidase, penicillin-G-amidase, β -lactamase, β -glucosidase, nitroreductase and carboxypeptidase A.

30 [0153] Techniques for conjugating therapeutic agents to proteins, and in particular to antibodies, are well-known. (*See, e.g.*, Arnon *et al.*, “Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy,” in *Monoclonal Antibodies And Cancer Therapy* (Reisfeld *et al.* eds., Alan R. Liss, Inc., 1985); Hellstrom *et al.*, “Antibodies For Drug Delivery,” in *Controlled Drug Delivery* (Robinson *et al.* eds., Marcel Dekker, Inc., 2nd
35 ed. 1987); Thorpe, “Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review,” in *Monoclonal Antibodies '84: Biological And Clinical Applications* (Pinchera *et al.* eds.,

- 5 1985); "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody In Cancer Therapy," in *Monoclonal Antibodies For Cancer Detection And Therapy* (Baldwin *et al.* eds., Academic Press, 1985); and Thorpe *et al.*, 1982, *Immunol. Rev.* 62:119-58. *See also, e.g.*, PCT publication WO 89/12624.)

[0154] In accordance with the methods described herein, the anti-CD70 ADC or ADC
10 derivative is internalized and accumulates within a CD70-expressing cell, where the ADC or ADC derivative exerts a therapeutic effect (*e.g.*, a cytotoxic, cytostatic, or immunosuppressive effect). Methods for determining accumulation and rates of accumulation are found in U.S. Provisional Application Serial No. 60/400,404, filed July 31, 2002; the disclosure of which is incorporated by reference herein.

15 [0155] Typically, when using an anti-CD70 antibody or derivative thereof conjugated to a therapeutic agent (*e.g.*, a drug or a prodrug converting enzyme), the agent is preferentially active when internalized by cells of the cancer to be treated or by activated immune cells (*e.g.*, activated lymphocytes or dendritic cells). In other embodiments, the anti-CD70 ADC or ADC derivative is not internalized, and the drug is effective to deplete or inhibiting CD70-
20 expressing cells by binding to the cell membrane. In yet other embodiments, anti-CD70 antibodies ADC or ADC derivatives thereof can be targeted to a biological molecules in a cell (*e.g.*, an inflammatory agent) and accumulate at or adjacent cells secreting or binding the biological molecule, where the therapeutic agent exerts an effect (*e.g.*, a cytotoxic, cytostatic, or immunosuppressive effect).

25 [0156] To minimize activity of the therapeutic agent outside the activated immune cells or CD70-expressing cancer cells, an antibody that specifically binds to cell membrane-bound CD70, but not to soluble CD70, can be used, so that the therapeutic agent is concentrated at the cell surface of the activated immune cell or CD70-expressing cancer cell. Alternatively, in a more typical embodiment, the therapeutic agent is conjugated in a manner that reduces its
30 activity unless it is cleaved off the antibody (*e.g.*, by hydrolysis or by a cleaving agent). In such embodiments, the therapeutic agent is attached to the antibody or derivative thereof with a cleavable linker that is sensitive to cleavage in the intracellular environment of the activated immune cell or CD70-expressing cancer cell but is not substantially sensitive to the extracellular environment, such that the conjugate is cleaved from the antibody or derivative
35 thereof when it is internalized by the activated immune cell or CD70-expressing cancer cell

- 5 (e.g., in the endosomal or, for example by virtue of pH sensitivity or protease sensitivity, in the lysosomal environment or in a caveolea). (See Section III(A), *infra*.)

[0157] Further, in certain embodiments, the ADC or ADC derivative comprises a therapeutic agent that is charged relative to the plasma membrane, thereby further minimizing the ability of the agent to cross the plasma membrane once internalized by a cell. As used herein, a
10 “charged agent” means an agent that (a) is polarized, such that one region of the agent has a charge relative to the plasma membrane, or (b) has a net charge relative to the plasma membrane.

[0158] Typically, the anti-CD70 antibody-drug conjugate (ADC) or ADC derivative is substantially purified (e.g., substantially free from substances that limit its effect or produce
15 undesired side-effects). In certain specific embodiments, the anti-CD70 ADC or ADC derivative is 40% pure, more typically about 50% pure, and most typically about 60% pure. In other specific embodiments, the anti-CD70 ADC or ADC derivative is at least approximately 60-65%, 65-70%, 70-75%, 75-80%, 80-85%, 85-90%, 90-95%, or 95-98% pure. In another specific embodiment, the anti-CD70 ADC or ADC derivative is
20 approximately 99% pure.

A. Linkers

[0159] Typically, the ADC or ADC derivative comprises a linker region between the therapeutic agent and the anti-CD70 antibody or derivative thereof. As noted *supra*, in
typical embodiments, the linker is cleavable under intracellular conditions, such that cleavage
25 of the linker releases the therapeutic agent from the antibody in the intracellular environment.

[0160] For example, in some embodiments, the linker is cleavable by a cleaving agent that is present in the intracellular environment (e.g., within a lysosome or endosome or caveolea). The linker can be, e.g., a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease.
30 Typically, the peptidyl linker is at least two amino acids long or at least three amino acids long. Cleaving agents can include cathepsins B and D and plasmin, all of which are known to hydrolyze dipeptide drug derivatives resulting in the release of active drug inside target cells (see, e.g., Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123). Most typical are peptidyl linkers that are cleavable by enzymes that are present in CD70-expressing cells.
35 For example, a peptidyl linker that is cleavable by the thiol-dependent protease cathepsin-B, which is highly expressed in cancerous tissue, can be used (e.g., a Phe-Leu or a Gly-Phe-Leu-

5 Gly linker). Other such linkers are described, *e.g.*, in U.S. Patent No. 6,214,345. In specific embodiments, the peptidyl linker cleavable by an intracellular protease is a Val-Cit linker or a Phe-Lys linker (*see, e.g.*, U.S. patent 6,214,345, which describes the synthesis of doxorubicin with the val-cit linker). One advantage of using intracellular proteolytic release of the therapeutic agent is that the agent is typically attenuated when conjugated and the serum
10 stabilities of the conjugates are typically high.

[0161] In other embodiments, the cleavable linker is pH-sensitive, *i.e.*, sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker hydrolyzable under acidic conditions. For example, an acid-labile linker that is hydrolyzable in the lysosome (*e.g.*, a hydrazone, semicarbazone, thiosemicarbazone, cis-aconitic amide, orthoester, acetal, ketal, or the like)
15 can be used. (*See, e.g.*, U.S. Patent Nos. 5,122,368; 5,824,805; 5,622,929; Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123; Neville *et al.*, 1989, *Biol. Chem.* 264:14653-14661.) Such linkers are relatively stable under neutral pH conditions, such as those in the blood, but are unstable at below pH 5.5 or 5.0, the approximate pH of the lysosome. In certain embodiments, the hydrolyzable linker is a thioether linker (such as, *e.g.*, a thioether
20 attached to the therapeutic agent via an acylhydrazone bond (*see, e.g.*, U.S. Patent No. 5,622,929)).

[0162] In yet other embodiments, the linker is cleavable under reducing conditions (*e.g.*, a disulfide linker). A variety of disulfide linkers are known in the art, including, for example, those that can be formed using SATA (N-succinimidyl-S-acetylthioacetate), SPDP (N-succinimidyl-3-(2-pyridyldithio)propionate), SPDB (N-succinimidyl-3-(2-pyridyldithio)butyrate) and SMPT (N-succinimidyl-oxycarbonyl-alpha-methyl-alpha-(2-pyridyl-dithio)toluene), SPDB and SMPT (*See, e.g.*, Thorpe *et al.*, 1987, *Cancer Res.* 47:5924-5931; Wawrzynczak *et al.*, In *Immunoconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer* (C. W. Vogel ed., Oxford U. Press, 1987. *See also*
25 U.S. Patent No. 4,880,935.)

[0163] In yet other specific embodiments, the linker is a malonate linker (Johnson *et al.*, 1995, *Anticancer Res.* 15:1387-93), a maleimidobenzoyl linker (Lau *et al.*, 1995, *Bioorg-Med-Chem.* 3(10):1299-1304), or a 3'-N-amide analog (Lau *et al.*, 1995, *Bioorg-Med-Chem.* 3(10):1305-12).

35 [0164] Typically, the linker is not substantially sensitive to the extracellular environment. As used herein, "not substantially sensitive to the extracellular environment," in the context of a

5 linker, means that no more than about 20%, typically no more than about 15%, more typically no more than about 10%, and even more typically no more than about 5%, no more than about 3%, or no more than about 1% of the linkers, in a sample of ADC or ADC derivative, are cleaved when the ADC or ADC derivative present in an extracellular environment (*e.g.*, in plasma). Whether a linker is not substantially sensitive to the extracellular environment
 10 can be determined, for example, by incubating independently with plasma both (a) the ADC or ADC derivative (the "ADC sample") and (b) an equal molar amount of unconjugated antibody or therapeutic agent (the "control sample") for a predetermined time period (*e.g.*, 2, 4, 8, 16, or 24 hours) and then comparing the amount of unconjugated antibody or therapeutic agent present in the ADC sample with that present in control sample, as measured, for
 15 example, by high performance liquid chromatography.

[0165] In other, non-mutually exclusive embodiments, the linker promotes cellular internalization. In certain embodiments, the linker promotes cellular internalization when conjugated to the therapeutic agent (*i.e.*, in the milieu of the linker-therapeutic agent moiety of the ADC or ADC derivative as described herein). In yet other embodiments, the linker
 20 promotes cellular internalization when conjugated to both the therapeutic agent and the anti-CD70 antibody or derivative thereof (*i.e.*, in the milieu of the ADC or ADC derivative as described herein).

[0166] A variety of linkers that can be used with the present compositions and methods are described in WO 2004010957 entitled "Drug Conjugates and Their Use for Treating Cancer,
 25 An Autoimmune Disease or an Infectious Disease" filed July 31, 2003, and U.S. Provisional Application No. 60/400,403, entitled "Drug Conjugates and their use for treating cancer, an autoimmune disease or an infectious disease", filed July 31, 2002 (the disclosure of which is incorporated by reference herein).

[0167] In certain embodiments, the linker unit has the following general formula:



wherein:

-T- is a stretcher unit;

a is 0 or 1;

each -W- is independently an amino acid unit;

5 w is independently an integer ranging from 2 to 12;


-Y- is a spacer unit; and

y is 0, 1 or 2.

1. The Stretcher Unit


[0168] The stretcher unit (-T-), when present, links the anti-CD70 antibody unit to an amino acid unit (-W-). Useful functional groups that can be present on an anti-CD70 antibody, either naturally or via chemical manipulation include, but are not limited to, sulfhydryl, amino, hydroxyl, the anomeric hydroxyl group of a carbohydrate, and carboxyl. Suitable functional groups are sulfhydryl and amino. Sulfhydryl groups can be generated by reduction of the intramolecular disulfide bonds of an anti-CD70 antibody. Alternatively, sulfhydryl groups can be generated by reaction of an amino group of a lysine moiety of an anti-CD70 antibody with 2-iminothiolane (Traut's reagent) or other sulfhydryl generating reagents. In specific embodiments, the anti-CD70 antibody is a recombinant antibody and is engineered to carry one or more lysines. In other embodiments, the recombinant anti-CD70 antibody is engineered to carry additional sulfhydryl groups, *e.g.*, additional cysteines.

20 [0169] In certain specific embodiments, the stretcher unit forms a bond with a sulfur atom of the anti-CD70 antibody unit. The sulfur atom can be derived from a sulfhydryl (-SH) group of a reduced anti-CD70 antibody (A). Representative stretcher units of these embodiments are depicted within the square brackets of Formulas (Ia) and (Ib; see *infra*), wherein A-, -W-, -Y-, -D, w and y are as defined above and R¹ is selected from -C₁-C₁₀ alkylene-, -C₃-C₈ carbocyclo-, -O-(C₁-C₈ alkyl)-, -arylene-, -C₁-C₁₀ alkylene-arylene-, -arylene-C₁-C₁₀ alkylene-, -C₁-C₁₀ alkylene-(C₃-C₈ carbocyclo)-, -(C₃-C₈ carbocyclo)-C₁-C₁₀ alkylene-, -C₃-C₈ heterocyclo-, -C₁-C₁₀ alkylene-(C₃-C₈ heterocyclo)-, -(C₃-C₈ heterocyclo)-C₁-C₁₀ alkylene-, -(CH₂CH₂O)_r, and -(CH₂CH₂O)_r-CH₂-; and r is an integer ranging from 1-10.




(Ia)

5




[0170] An illustrative stretcher unit is that of formula (Ia) where R^1 is $-(\text{CH}_2)_5-$:




10

[0171] Another illustrative stretcher unit is that of formula (Ia) where R^1 is $-(\text{CH}_2\text{CH}_2\text{O})_r-$ CH_2- and r is 2:




[0172] Still another illustrative stretcher unit is that of formula (Ib) where R^1 is $-(\text{CH}_2)_5-$:



15

[0173] In certain other specific embodiments, the stretcher unit is linked to the anti-CD70 antibody unit (A) via a disulfide bond between a sulfur atom of the anti-CD70 antibody unit and a sulfur atom of the stretcher unit. A representative stretcher unit of this embodiment is depicted within the square brackets of Formula (II), wherein R^1 , A-, -W-, -Y-, -D, w and y are as defined above.




20

(II)

[0174] In other specific embodiments, the reactive group of the stretcher contains a reactive site that can be reactive to an amino group of an anti-CD70 antibody. The amino group can be that of an arginine or a lysine. Suitable amine reactive sites include, but are not limited to,


- 5 activated esters such as succinimide esters, 4-nitrophenyl esters, pentafluorophenyl esters, anhydrides, acid chlorides, sulfonyl chlorides, isocyanates and isothiocyanates.

Representative stretcher units of these embodiments are depicted within the square brackets of Formulas (IIIa) and (IIIb), wherein R^1 , A-, -W-, -Y-, -D, w and y are as defined above;




10 (IIIa)

(IIIb)



- 15 [0175] In yet another aspect, the reactive function of the stretcher contains a reactive site that is reactive to a modified carbohydrate group that can be present on an anti-CD70 antibody.


In a specific embodiment, the anti-CD70 antibody is glycosylated enzymatically to provide a carbohydrate moiety. The carbohydrate may be mildly oxidized with a reagent such as sodium periodate and the resulting carbonyl unit of the oxidized carbohydrate can be
 20 condensed with a stretcher that contains a functionality such as a hydrazide, an oxime, a reactive amine, a hydrazine, a thiosemicarbazide, a hydrazine carboxylate, and an arylhydrazide such as those described by Kaneko *et al.*, 1991, *Bioconjugate Chem* 2:133-41. Representative stretcher units of this embodiment are depicted within the square brackets of Formulas (IVa)-(IVc), wherein R^1 , A-, -W-, -Y-, -D, w and y are as defined above.



25


(IVa)

5



(IVb)

10




(IVc)

2. The Amino Acid Unit

[0176] The amino acid unit (-W-) links the stretcher unit (-T-) to the Spacer unit (-Y-) if the Spacer unit is present, and links the stretcher unit to the cytotoxic or cytostatic agent (Drug unit; D) if the spacer unit is absent.


[0177] -W_w- is a dipeptide, tripeptide, tetrapeptide, pentapeptide, hexapeptide, heptapeptide, octapeptide, nonapeptide, decapeptide, undecapeptide or dodecapeptide unit. Each -W- unit independently has the formula denoted below in the square brackets, and w is an integer ranging from 2 to 12:

20




25

wherein R² is hydrogen, methyl, isopropyl, isobutyl, sec-butyl, benzyl, p-hydroxybenzyl, -CH₂OH, -CH(OH)CH₃, -CH₂CH₂SCH₃, -CH₂CONH₂, -CH₂COOH, -CH₂CH₂CONH₂, -CH₂CH₂COOH, -(CH₂)₃NHC(=NH)NH₂, -(CH₂)₃NH₂, -(CH₂)₃NHCOCH₃, -(CH₂)₃NHCHO, -(CH₂)₄NHC(=NH)NH₂, -(CH₂)₄NH₂, -(CH₂)₄NHCOCH₃, -(CH₂)₄NHCHO, -(CH₂)₃NHCONH₂, -(CH₂)₄NHCONH₂, -CH₂CH₂CH(OH)CH₂NH₂, 2-pyridylmethyl-, 3-pyridylmethyl-, 4-pyridylmethyl-, phenyl, cyclohexyl,



[0178] The amino acid unit of the linker unit can be enzymatically cleaved by an enzyme including, but not limited to, a tumor-associated protease to liberate the drug unit (-D) which is protonated *in vivo* upon release to provide a cytotoxic drug (D).


[0179] Illustrative W_w units are represented by formulas (V)-(VII):




(V)

wherein R^3 and R^4 are as follows:

R^3	R^4
Benzyl	$(CH_2)_4NH_2$;
Methyl	$(CH_2)_4NH_2$;
Isopropyl	$(CH_2)_4NH_2$;
Isopropyl	$(CH_2)_3NHCONH_2$;
Benzyl	$(CH_2)_3NHCONH_2$;



5




(VI)

wherein R^3 , R^4 and R^5 are as follows:

R^3	R^4	R^5
Benzyl	benzyl	$(\text{CH}_2)_4\text{NH}_2$;
Isopropyl	benzyl	$(\text{CH}_2)_4\text{NH}_2$; and
H	benzyl	$(\text{CH}_2)_4\text{NH}_2$;

10



(VII)

wherein R^3 , R^4 , R^5 and R^6 are as follows:

15

R^3	R^4	R^5	R^6
H	Benzyl	isobutyl	H; and
methyl	Isobutyl	methyl	isobutyl.

5 [0180] Suitable amino acid units include, but are not limited to, units of formula (V) where: R^3 is benzyl and R^4 is $-(CH_2)_4NH_2$; R^3 is isopropyl and R^4 is $-(CH_2)_4NH_2$; or R^3 is isopropyl and R^4 is $-(CH_2)_3NHCONH_2$. Another suitable amino acid unit is a unit of formula (VI), where: R^3 is benzyl, R^4 is benzyl, and R^5 is $-(CH_2)_4NH_2$.

[0181] $-W_w-$ units can be designed and optimized in their selectivity for enzymatic cleavage
10 by a particular tumor-associated protease. The suitable $-W_w-$ units are those whose cleavage is catalyzed by the proteases, cathepsin B, C and D, and plasmin.

[0182] In one embodiment, $-W_w-$ is a dipeptide, tripeptide or tetrapeptide unit.

[0183] Where R^2 , R^3 , R^4 , R^5 or R^6 is other than hydrogen, the carbon atom to which R^2 , R^3 , R^4 , R^5 or R^6 is attached is chiral.

15 [0184] Each carbon atom to which R^2 , R^3 , R^4 , R^5 or R^6 is attached is independently in the (S) or (R) configuration.

[0185] In a certain embodiment, the amino acid unit is a phenylalanine-lysine dipeptide (Phe-Lys or FK linker). In another embodiment, the amino acid unit is a valine-citrulline dipeptide (Val-Cit or VC linker).


20 3. The Spacer Unit

[0186] The spacer unit ($-Y-$), when present, links an amino acid unit to the drug unit. Spacer units are of two general types: self-immolative and non self-immolative. A non self-immolative spacer unit is one in which part or all of the spacer unit remains bound to the drug unit after enzymatic cleavage of an amino acid unit from the anti-CD70 antibody-linker-drug
25 conjugate or the drug-linker compound. Examples of a non self-immolative spacer unit include, but are not limited to a (glycine-glycine) spacer unit and a glycine spacer unit (both depicted in Scheme 1). When an anti-CD70 antibody-linker-drug conjugate containing a glycine-glycine spacer unit or a glycine spacer unit undergoes enzymatic cleavage via a tumor-cell associated-protease, a cancer-cell-associated protease or a lymphocyte-associated
30 protease, a glycine-glycine-drug moiety or a glycine-drug moiety is cleaved from A-T- W_w- . To liberate the drug, an independent hydrolysis reaction should take place within the target cell to cleave the glycine-drug unit bond.

- 5 [0187] In a typical embodiment, -Y- is a *p*-aminobenzyl ether which can be substituted with Q_m where Q is is -C₁-C₈ alkyl, -C₁-C₈ alkoxy, -halogen, - nitro or -cyano; and m is an integer ranging from 0-4.

Scheme 1

[0188] In one embodiment, a non self-immolative spacer unit (-Y-) is -Gly-Gly-.



[0189] In another embodiment, a non self-immolative the spacer unit (-Y-) is -Gly-.


[0190] In one embodiment, the drug-linker compound or an anti-CD70 antibody-linker-drug conjugate lacks a spacer unit (y=0).

- 15 [0191] Alternatively, an anti-CD70 antibody-linker-drug conjugate containing a self-immolative spacer unit can release the drug (D) without the need for a separate hydrolysis step. In these embodiments, -Y- is a *p*-aminobenzyl alcohol (PAB) unit that is linked to -W_w- via the nitrogen atom of the PAB group, and connected directly to -D via a carbonate, carbamate or ether group (Scheme 2 and Scheme 3).

20


25

5

Scheme 2

where Q is -C₁-C₈ alkyl, -C₁-C₈ alkoxy, -halogen, -nitro or -cyano; m is an integer ranging from 0-4; and p is an integer ranging from 1-20.

5


Scheme 3

where Q is -C₁-C₈ alkyl, -C₁-C₈ alkoxy, -halogen, -nitro or -cyano; m is an integer ranging from 0-4; and p is an integer ranging from 1-20.

[0192] Other examples of self-immolative spacers include, but are not limited to, aromatic compounds that are electronically equivalent to the PAB group such as 2-aminoimidazol-5-methanol derivatives (*see Hay et al.*, 1999, *Bioorg. Med. Chem. Lett.* 9:2237 for examples) and ortho or para-aminobenzylacetals. Spacers can be used that undergo facile cyclization upon amide bond hydrolysis, such as substituted and unsubstituted 4-aminobutyric acid amides (Rodrigues *et al.*, 1995, *Chemistry Biology* 2:223), appropriately substituted bicyclo[2.2.1] and bicyclo[2.2.2] ring systems (Storm *et al.*, 1972, *J. Amer. Chem. Soc.* 94:5815) and 2-aminophenylpropionic acid amides (Amsberry *et al.*, 1990, *J. Org. Chem.* 55:5867). Elimination of amine-containing drugs that are substituted at the α-position of glycine (Kingsbury, *et al.*, 1984, *J. Med. Chem.* 27:1447) are also examples of self-immolative spacer strategies that can be applied to the anti-CD70 antibody-linker-drug conjugates.

[0193] In an alternate embodiment, the spacer unit is a branched bis(hydroxymethyl)styrene (BHMS) unit (Scheme 4), which can be used to incorporate additional drugs.

5


Scheme 4

where Q is -C₁-C₈ alkyl, -C₁-C₈ alkoxy, -halogen, -nitro or -cyano; m is an integer ranging from 0-4; n is 0 or 1; and p is an integer ranging from 1-20.

10 [0194] In one embodiment, the two -D moieties are the same.


[0195] In another embodiment, the two -D moieties are different.

[0196] Typical spacer units (-Y_y-) are represented by Formulas (VIII)-(X):




(VIII)

15 where Q is C₁-C₈ alkyl, C₁-C₈ alkoxy, halogen, nitro or cyano; and m is an integer ranging from 0-4;



(IX); and



20

(X)

5 B. Therapeutic Agents

- [0197] In accordance with the methods described herein, any agent that exerts a therapeutic effect on cancer cells or activated immune cells can be used as the therapeutic agent for conjugation to an anti-CD70 antibody or derivative thereof. (*See, e.g.*, WO 2004/010957, "Drug Conjugates and Their Use for Treating Cancer, An Autoimmune Disease or an Infectious Disease" (*supra*) and U.S. Provisional Application No. 60/400,403 (*supra*)). Typically, the therapeutic agent is a cytotoxic or immunosuppressive agent.
- [0198] Useful classes of cytotoxic or immunosuppressive agents include, for example, antitubulin agents, auristatins, DNA minor groove binders, DNA replication inhibitors, alkylating agents (*e.g.*, platinum complexes such as *cis*-platin, mono(platinum), bis(platinum) and tri-nuclear platinum complexes and carboplatin), anthracyclines, antibiotics, antifolates, antimetabolites, chemotherapy sensitizers, duocarmycins, etoposides, fluorinated pyrimidines, ionophores, lexitropsins, nitrosoureas, platinols, pre-forming compounds, purine antimetabolites, puromycins, radiation sensitizers, steroids, taxanes, topoisomerase inhibitors, vinca alkaloids, or the like.
- [0199] Individual cytotoxic or immunosuppressive agents include, for example, an androgen, anthramycin (AMC), asparaginase, 5-azacytidine, azathioprine, bleomycin, busulfan, buthionine sulfoximine, camptothecin, carboplatin, carmustine (BSNU), CC-1065, chlorambucil, cisplatin, colchicine, cyclophosphamide, cytarabine, cytidine arabinoside, cytochalasin B, dacarbazine, dactinomycin (formerly actinomycin), daunorubicin, decarbazine, docetaxel, doxorubicin, an estrogen, 5-fluorodeoxyuridine, 5-fluorouracil, gramicidin D, hydroxyurea, idarubicin, ifosfamide, irinotecan, lomustine (CCNU), mechlorethamine, melphalan, 6-mercaptopurine, methotrexate, mithramycin, mitomycin C, mitoxantrone, nitroimidazole, paclitaxel, plicamycin, procarbazine, streptozotocin, tenoposide, 6-thioguanine, thioTEPA, topotecan, vinblastine, vincristine, vinorelbine, VP-16 and VM-26.
- [0200] In some typical embodiments, the therapeutic agent is a cytotoxic agent. Suitable cytotoxic agents include, for example, dolastatins (*e.g.*, auristatin E, AFP, MMAF, MMAE), DNA minor groove binders (*e.g.*, enediynes and lexitropsins), duocarmycins, taxanes (*e.g.*, paclitaxel and docetaxel), puromycins, vinca alkaloids, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, echinomycin,

5 combretastatin, netropsin, epothilone A and B, estramustine, cryptophysins, cemadotin, maytansinoids, discodermolide, eleutherobin, and mitoxantrone.

[0201] In certain embodiments, the cytotoxic agent is a conventional chemotherapeutic such as, for example, doxorubicin, paclitaxel, melphalan, vinca alkaloids, methotrexate, mitomycin C or etoposide. In addition, potent agents such as CC-1065 analogues, calicheamicin,
10 maytansine, analogues of dolastatin 10, rhizoxin, and palytoxin can be linked to the anti-CD70 antibodies or derivatives thereof.

[0202] In specific embodiments, the cytotoxic or cytostatic agent is auristatin E (also known in the art as dolastatin-10) or a derivative thereof. Typically, the auristatin E derivative is, *e.g.*, an ester formed between auristatin E and a keto acid. For example, auristatin E can be
15 reacted with paraacetyl benzoic acid or benzoylvaleric acid to produce AEB and AEVB, respectively. Other typical auristatin derivatives include AFP, MMAF, and MMAE. The synthesis and structure of auristatin E and its derivatives are described in U.S. Patent Application Nos. 09/845,786 (U.S. Patent Application Publication No. 20030083263) and 10/001,191; International Patent Application No. PCT/US03/24209, International Patent
20 Application No. PCT/US02/13435, and U.S. Patent Nos. 6,323,315; 6,239,104; 6,034,065; 5,780,588; 5,665,860; 5,663,149; 5,635,483; 5,599,902; 5,554,725; 5,530,097; 5,521,284; 5,504,191; 5,410,024; 5,138,036; 5,076,973; 4,986,988; 4,978,744; 4,879,278; 4,816,444; and 4,486,414.

[0203] In specific embodiments, the cytotoxic agent is a DNA minor groove binding agent.
25 (*See, e.g.*, U.S. Patent No. 6,130,237.) For example, in certain embodiments, the minor groove binding agent is a CBI compound. In other embodiments, the minor groove binding agent is an enediyne (*e.g.*, calicheamicin).

[0204] In certain embodiments, the ADC or ADC derivative comprises an anti-tubulin agent. Examples of anti-tubulin agents include, but are not limited to, taxanes (*e.g.*, Taxol®
30 (paclitaxel), Taxotere® (docetaxel)), T67 (Tularik), vinca alkyls (*e.g.*, vincristine, vinblastine, vindesine, and vinorelbine), and dolastatins (*e.g.*, auristatin E, AFP, MMAF, MMAE, AEB, AEVB). Other antitubulin agents include, for example, baccatin derivatives, taxane analogs (*e.g.*, epothilone A and B), nocodazole, colchicine and colcimid, estramustine, cryptophysins, cemadotin, maytansinoids, combretastatins, discodermolide, and eleutherobin.

5 [0205] In certain embodiments, the cytotoxic agent is a maytansinoid, another group of anti-tubulin agents. For example, in specific embodiments, the maytansinoid is maytansine or DM-1 (ImmunoGen, Inc.; *see also* Chari *et al.*, 1992, *Cancer Res.* 52:127-131).

[0206] In certain embodiments, the therapeutic agent is not a radioisotope.

10 [0207] In certain embodiments, the cytotoxic or immunosuppressive agent is an antimetabolite. The antimetabolite can be, for example, a purine antagonist (*e.g.* azothioprine or mycophenolate mofetil), a dihydrofolate reductase inhibitor (*e.g.*, methotrexate), acyclovir, gangcyclovir, zidovudine, vidarabine, ribavirin, azidothymidine, cytidine arabinoside, amantadine, dideoxyuridine, iododeoxyuridine, poscarnet, or trifluridine.

[0208] In other embodiments, the cytotoxic or immunosuppressive agent is tacrolimus, 15 cyclosporine or rapamycin. In further embodiments, the cytotoxic agent is aldesleukin, alemtuzumab, alitretinoin, allopurinol, altretamine, amifostine, anastrozole, arsenic trioxide, bexarotene, bexarotene, calusterone, capecitabine, celecoxib, cladribine, Darbepoetin alfa, Denileukin diftitox, dexrazoxane, dromostanolone propionate, epirubicin, Epoetin alfa, estramustine, exemestane, Filgrastim, floxuridine, fludarabine, fulvestrant, gemcitabine, 20 gemtuzumab ozogamicin, goserelin, idarubicin, ifosfamide, imatinib mesylate, Interferon alfa-2a, irinotecan, letrozole, leucovorin, levamisole, meclorethamine or nitrogen mustard, megestrol, mesna, methotrexate, methoxsalen, mitomycin C, mitotane, nandrolone phenpropionate, oprelvekin, oxaliplatin, pamidronate, pegademase, pegaspargase, pegfilgrastim, pentostatin, pipobroman, plicamycin, porfimer sodium, procarbazine, 25 quinacrine, rasburicase, Rituximab, Sargramostim, streptozocin, tamoxifen, temozolomide, teniposide, testolactone, thioguanine, toremifene, Tositumomab, Trastuzumab, tretinoin, uracil mustard, valrubicin, vinblastine, vincristine, vinorelbine and zoledronate.

[0209] In additional embodiments, the drug is a humanized anti HER2 monoclonal antibody, RITUXAN (rituximab; Genentech; a chimeric anti CD20 monoclonal antibody); 30 OVAREX (AltaRex Corporation, MA); PANOREX (Glaxo Wellcome, NC; a murine IgG2a antibody); Cetuximab Erbitux (Imclone Systems Inc., NY; an anti-EGFR IgG chimeric antibody); Vitaxin (MedImmune, Inc., MD; Campath I/H (Leukosite, MA; a humanized IgG1 antibody); Smart MI95 (Protein Design Labs, Inc., CA; a humanized anti-CD33 IgG antibody); LymphoCide (Immunomedics, Inc., NJ; a humanized anti-CD22 IgG antibody); 35 Smart ID10 (Protein Design Labs, Inc., CA; a humanized anti-HLA-DR antibody); Oncolym (Techniclone, Inc., CA; a radiolabeled murine anti-HLA-Dr10 antibody); Allomune

- 5 (BioTransplant, CA; a humanized anti-CD2 mAb); Avastin (Genentech, Inc., CA; an anti-VEGF humanized antibody); Epratuzamab (Immunomedics, Inc., NJ and Amgen, CA; an anti-CD22 antibody); and CEAcide (Immunomedics, NJ; a humanized anti-CEA antibody).

[0210] Other suitable antibodies include, but are not limited to, antibodies against the following antigens: CA125, CA15-3, CA19-9, L6, Lewis Y, Lewis X, alpha fetoprotein, CA
10 242, placental alkaline phosphatase, prostate specific antigen, prostatic acid phosphatase, epidermal growth factor, MAGE-1, MAGE-2, MAGE-3, MAGE -4, anti transferrin receptor, p97, MUC1-KLH, CEA, gp100, MART1, Prostate Specific Antigen, IL-2 receptor, CD20, CD52, CD33, CD22, human chorionic gonadotropin, CD38, CD40, mucin, P21, MPG, and Neu oncogene product.

- 15 [0211] In certain embodiments, the therapeutic agent is an immunosuppressive agent. The immunosuppressive agent can be, for example, gancyclovir, etanercept, tacrolimus, cyclosporine, rapamycin, cyclophosphamide, azathioprine, mycophenolate mofetil or methotrexate. Alternatively, the immunosuppressive agent can be, for example, a glucocorticoid (*e.g.*, cortisol or aldosterone) or a glucocorticoid analogue (*e.g.*, prednisone or
20 dexamethasone).

[0212] In certain typical embodiments, the immunosuppressive agent is an anti-inflammatory agent, such as arylcarboxylic derivatives, pyrazole-containing derivatives, oxycam derivatives and nicotinic acid derivatives. Classes of anti-inflammatory agents include, for example, cyclooxygenase inhibitors, 5-lipoxygenase inhibitors, and leukotriene receptor antagonists.

- 25 [0213] Suitable cyclooxygenase inhibitors include meclofenamic acid, mefenamic acid, carprofen, diclofenac, diflunisal, fenbufen, fenoprofen, ibuprofen, indomethacin, ketoprofen, nabumetone, naproxen, sulindac, tenoxicam, tolmetin, and acetylsalicylic acid.



- [0214] Suitable lipoxygenase inhibitors include redox inhibitors (*e.g.*, catechol butane derivatives, nordihydroguaiaretic acid (NDGA), masoprocol, phenidone, lanopalen,
30 indazolinones, naphazatrom, benzofuranol, alkylhydroxylamine), and non-redox inhibitors (*e.g.*, hydroxythiazoles, methoxyalkylthiazoles, benzopyrans and derivatives thereof, methoxytetrahydropyran, boswellic acids and acetylated derivatives of boswellic acids, and quinolinemethoxyphenylacetic acids substituted with cycloalkyl radicals), and precursors of redox inhibitors.

- 5 [0215] Other suitable lipoxygenase inhibitors include antioxidants (*e.g.*, phenols, propyl gallate, flavonoids and/or naturally occurring substrates containing flavonoids, hydroxylated derivatives of the flavones, flavonol, dihydroquercetin, luteolin, galangin, orobol, derivatives of chalcone, 4,2',4'-trihydroxychalcone, ortho-aminophenols, N-hydroxyureas, benzofuranols, ebselen and species that increase the activity of the reducing selenoenzymes),
- 10 iron chelating agents (*e.g.*, hydroxamic acids and derivatives thereof, N-hydroxyureas, 2-benzyl-1-naphthol, catechols, hydroxylamines, carnosol trolox C, catechol, naphthol, sulfasalazine, zileuton, 5-hydroxyanthranilic acid and 4-(omega-arylalkyl)phenylalkanoic acids), imidazole-containing compounds (*e.g.*, ketoconazole and itraconazole), phenothiazines, and benzopyran derivatives.
- 15 [0216] Yet other suitable lipoxygenase inhibitors include inhibitors of eicosanoids (*e.g.*, octadecatetraenoic, eicosatetraenoic, docosapentaenoic, eicosahexaenoic and docosahexaenoic acids and esters thereof, PGE1 (prostaglandin E1), PGA2 (prostaglandin A2), viprostol, 15-monohydroxyeicosatetraenoic, 15-monohydroxy-eicosatrienoic and 15-monohydroxyeicosapentaenoic acids, and leukotrienes B5, C5 and D5), compounds
- 20 interfering with calcium flows, phenothiazines, diphenylbutylamines, verapamil, fuscoidin, curcumin, chlorogenic acid, caffeic acid, 5,8,11,14-eicosatetraenoic acid (ETYA), hydroxyphenylretinamide, lonapalen, esculin, diethylcarbamazine, phenantrolin, baicalein, proxidromil, thioethers, diallyl sulfide and di-(1-propenyl) sulfide.
- [0217] Leukotriene receptor antagonists include calcitriol, ontazolast, Bayer Bay-x-1005,
- 25 Ciba-Geigy CGS-25019C, ebselen, Leo Denmark ETH-615, Lilly LY-293111, Ono ONO-4057, Terumo TMK-688, Boehringer Ingelheim BI-RM-270, Lilly LY 213024, Lilly LY 264086, Lilly LY 292728, Ono ONO LB457, Pfizer 105696, Perdue Frederick PF 10042, Rhone-Poulenc Rorer RP 66153, SmithKline Beecham SB-201146, SmithKline Beecham SB-201993, SmithKline Beecham SB-209247, Searle SC-53228, Sumitomo SM 15178,
- 30 American Home Products WAY 121006, Bayer Bay-o-8276, Warner-Lambert CI-987, Warner-Lambert CI-987BPC-15LY 223982, Lilly LY 233569, Lilly LY-255283, MacroNex MNX-160, Merck and Co. MK-591, Merck and Co. MK-886, Ono ONO-LB-448, Purdue Frederick PF-5901, Rhone-Poulenc Rorer RG 14893, Rhone-Poulenc Rorer RP 66364, Rhone-Poulenc Rorer RP 69698, Shionogi S-2474, Searle SC-41930, Searle SC-50505,
- 35 Searle SC-51146, Searle SC-52798, SmithKline Beecham SK&F-104493, Leo Denmark SR-2566, Tanabe T-757 and Teijin TEI-1338.

5

1. Dolastatin Drugs


[0218] In certain embodiments, the cytotoxic or cytostatic agent is a dolastatin. In more specific embodiments, the dolastatin is of the auristatin class. As used herein, the term dolastatin encompasses naturally occurring auristatins and non-naturally occurring derivatives, for example MMAE. Thus, in a specific embodiment, the cytotoxic or cytostatic agent is MMAE (Formula XI). In another specific embodiment, the cytotoxic or cytostatic agent is AFP (Formula XVI).



(XI)


15

[0219] In certain embodiments, the cytotoxic or cytostatic agent is a dolastatin of formulas XII-XXI.




(XII)

20




5

(XIII)




10


(XIV)



(XV)




15




5

(XVI)




(XVII)




10

(XVIII)




(XIV)



15

(XX)

5



(XXI)

C. Formation of Anti-CD70 ADCs and ADC Derivatives

10 [0220] The generation of anti-CD70 ADCs and ADC derivatives can be accomplished by any technique known to the skilled artisan. Briefly, the anti-CD70 ADCs comprise an anti-CD70 antibody, a drug, and optionally a linker that joins the drug and the antibody. A number of different reactions are available for covalent attachment of drugs to antibodies. This is often accomplished by reaction of the amino acid residues of the antibody molecule, including the

15 amine groups of lysine, the free carboxylic acid groups of glutamic and aspartic acid, the sulfhydryl groups of cysteine and the various moieties of the aromatic amino acids. One of the most commonly used non-specific methods of covalent attachment is the carbodiimide reaction to link a carboxy (or amino) group of a compound to amino (or carboxy) groups of the antibody. Additionally, bifunctional agents such as dialdehydes or imidoesters have been

20 used to link the amino group of a compound to amino groups of the antibody molecule. Also available for attachment of drugs to antibodies is the Schiff base reaction. This method involves the periodate oxidation of a drug that contains glycol or hydroxy groups, thus forming an aldehyde which is then reacted with the antibody molecule. Attachment occurs via formation of a Schiff base with amino groups of the antibody molecule. Isothiocyanates

25 can also be used as coupling agents for covalently attaching drugs to antibodies. Other techniques are known to the skilled artisan and within the scope of the present invention. Non-limiting examples of such techniques are described in, *e.g.*, U.S. Patent Nos. 5,665,358; 5,643,573; and 5,556,623, which are incorporated by reference in their entireties herein.

5 [0221] In certain embodiments, an intermediate, which is the precursor of the linker, is reacted with the drug under appropriate conditions. In certain embodiments, reactive groups are used on the drug and/or the intermediate. The product of the reaction between the drug and the intermediate, or the derivatized drug, is subsequently reacted with the anti-CD70 antibody under appropriate conditions.

10

IV. Other CD70-Targeting Moieties and Protein-Drug Conjugates

[0222] As indicated *supra*, in other embodiments, the CD70-targeting moiety need not be an antibody to be useful in accordance with the methods described herein. Accordingly, a CD70-targeting moiety can include one or more CDRs from an antibody that binds to CD70 and depletes or inhibits the proliferation of CD70-expressing cells when conjugated to a cytotoxic agent, or from an antibody that binds to CD70 and exerts an immunosuppressive effect when conjugated to an immunosuppressive agent. Typically, the protein is a multimer, most typically a dimer.

20 [0223] Further, CD70-binding proteins useful in accordance with the methods provided herein include fusion proteins, *i.e.*, proteins that are recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugation) to heterologous proteins (of typically at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or at least 100 amino acids). The fusion does not necessarily need to be direct, but may occur through linker sequences.

25 [0224] For example, CD70-targeting moieties useful in the present methods can be produced recombinantly by fusing the coding region of one or more of the CDRs of an anti-CD70 antibody in frame with a sequence coding for a heterologous protein. The heterologous protein may provide one or more of the following characteristics: added therapeutic benefits; promote stable expression; provide a means of facilitating high yield recombinant expression; and/or provide a multimerization domain.

30 [0225] In other aspects, CD70-targeting moieties can include CD27 and variants or fragments thereof that bind to CD70. CD70-targeting moieties can further include peptides, ligands and other molecules that specifically bind to CD70.

[0226] CD70-targeting moieties useful in the methods described herein can be identified using any method suitable for screening for protein-protein interactions. Typically, proteins are initially identified by their ability to specifically bind to CD70, then their ability to exert a

35

5 cytostatic or cytotoxic effect on activated lymphocytes or CD70-expressing cancer cells when
conjugated to a cytotoxic or cytostatic agent, or an immunosuppressive effect on an immune
cell when conjugated to an immunosuppressive agent. Among the traditional methods which
can be employed are "interaction cloning" techniques which entail probing expression
libraries with labeled CD70 in a manner similar to the technique of antibody probing of ?gt11
10 libraries. By way of example and not limitation, this can be achieved as follows: a cDNA
clone encoding CD70 (or a 1F6 or 2F2 binding domain thereof) is modified at the terminus
by inserting the phosphorylation site for the heart muscle kinase (HMK) (*see, e.g.,* Blanar and
Rutter, 1992, *Science* 256:1014-1018). The recombinant protein is expressed in *E. coli* and
purified on a GDP-affinity column to homogeneity (Edery *et al.*, 1988, *Gene* 74:517-525) and
15 labeled using $\gamma^{32}\text{P}$ -ATP and bovine heart muscle kinase (Sigma) to a specific activity of
 1×10^8 cpm/ μg , and used to screen a human placenta ?gt11 cDNA library in a "far-Western
assay" (Blanar and Rutter, 1992, *Science* 256:1014-1018). Plaques which interact with the
CD70 probe are isolated. The cDNA inserts of positive ? plaques are released and subcloned
into a vector suitable for sequencing, such as pBluescript KS (Stratagene).

20 [0227] One method which detects protein interactions *in vivo*, the two-hybrid system, is
described in detail for illustration purposes only and not by way of limitation. One version of
this system has been described (Chien *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88:9578-9582)
and is commercially available from Clontech (Palo Alto, CA).

[0228] Once a CD70-binding protein is identified, its ability (alone or when multimerized or
25 fused to a dimerization or multimerization domain) to exert a cytostatic or cytotoxic effect on
CD70-expressing cancer cells (when conjugated to a cytotoxic agent) or an
immunosuppressive effect on a CD70-expressing immune cell (when conjugated to an
immunosuppressive agent) is determined by the methods described *infra*.

30 V. Assays for Cytotoxic, Cytostatic, and Immunosuppressive Activities

[0229] In accordance with the methods described herein, an anti-CD70 ADC or ADC
derivative exerts a cytotoxic or cytostatic effect on a CD70-expressing cancer cell, or a
cytotoxic, cytostatic, or immunosuppressive effect on activated CD70-expressing immune
cell (*e.g.,* activated lymphocyte or dendritic cell). Activated lymphocytes that can be assayed
35 for a cytotoxic, cytostatic, or immunosuppressive effect of an anti-CD70 ADC or ADC
derivative can be cultured cell lines (*e.g.,* CESS, which is available from the ATCC; or KMH2

5 and L428, both of which are available from Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH), or from lymphocytes prepared from a fresh blood sample or other sources. Lymphocytes can be activated by the appropriate cocktails of antibodies and cytokines. For example, T lymphocytes can be activated using allogeneic stimulator cells or antigen-pulsed autologous antigen presenting cells and T cell clones can be activated by PHA
10 and cytokines in the presence of feeder cells, as described in Example 9, *infra*. CD70-expressing cancer cells that can be assayed for a cytotoxic or cytotoxic effect can be culture cell lines as described in Examples 7, *infra*, or from CD70-expressing cancer cells prepared from a subject.

[0230] Once an anti-CD70 ADC or ADC derivative is confirmed as exerting a cytotoxic or
15 cytostatic on CD70-expressing cancer cells or a cytotoxic, cytostatic, or immunosuppressive effect on activated immune cells, its therapeutic value can be validated in an animal model. Exemplary animal models of immunological disorders or CD70-expressing cancers are described in Section VI, *infra*.

[0231] Methods of determining whether an agent exerts a cytostatic or cytotoxic effect on a
20 cell are known. Illustrative examples of such methods are described *infra*.

[0232] For determining whether an anti-CD70 ADC or ADC derivative exerts a cytostatic effect on activated immune cells or CD70-expressing cancer cells, a thymidine incorporation assay may be used. For example, activated immune cells (*e.g.*, activated lymphocytes) or CD70-expressing cancer cells at a density of 5,000 cells/well of a 96-well plated can be
25 cultured for a 72-hour period and exposed to 0.5 μ Ci of 3 H-thymidine during the final 8 hours of the 72-hour period, and the incorporation of 3 H-thymidine into cells of the culture is measured in the presence and absence of the ADC or ADC derivative.

[0233] For determining cytotoxicity, necrosis or apoptosis (programmed cell death) can be measured. Necrosis is typically accompanied by increased permeability of the plasma
30 membrane; swelling of the cell, and rupture of the plasma membrane. Apoptosis is typically characterized by membrane blebbing, condensation of cytoplasm, and the activation of endogenous endonucleases. Determination of any of these effects on activated immune cells or CD70 expressing cancer cells indicates that an anti-CD70 ADC or ADC derivative is useful in the treatment or prevention of immunological disorders and CD70-expressing
35 cancers.

5 [0234] Cell viability can be measured by determining in a cell the uptake of a dye such as neutral red, trypan blue, or ALAMAR™ blue (*see, e.g., Page et al., 1993, Intl. J. of Oncology* 3:473-476). In such an assay, the cells are incubated in media containing the dye, the cells are washed, and the remaining dye, reflecting cellular uptake of the dye, is measured spectrophotometrically. The protein-binding dye sulforhodamine B (SRB) can also be used
10 to measure cytotoxicity (Skehan *et al.*, 1990, *J. Nat'l Cancer Inst.* 82:1107-12).

[0235] Alternatively, a tetrazolium salt, such as MTT, is used in a quantitative colorimetric assay for mammalian cell survival and proliferation by detecting living, but not dead, cells (*see, e.g., Mosmann, 1983, J. Immunol. Methods* 65:55-63).

[0236] Apoptosis can be quantitated by measuring, for example, DNA fragmentation.
15 Commercial photometric methods for the quantitative *in vitro* determination of DNA fragmentation are available. Examples of such assays, including TUNEL (which detects incorporation of labeled nucleotides in fragmented DNA) and ELISA-based assays, are described in *Biochemica*, 1999, no. 2, pp. 34-37 (Roche Molecular Biochemicals).

[0237] Apoptosis can also be determined by measuring morphological changes in a cell. For
20 example, as with necrosis, loss of plasma membrane integrity can be determined by measuring uptake of certain dyes (*e.g., a fluorescent dye such as, for example, acridine orange or ethidium bromide*). A method for measuring apoptotic cell number has previously been described by Duke and Cohen, *Current Protocols In Immunology* (Coligan *et al.* eds., 1992, pp. 3.17.1-3.17.16). Cells can be also labeled with a DNA dye (*e.g., acridine orange,*
25 *ethidium bromide, or propidium iodide*) and the cells observed for chromatin condensation and margination along the inner nuclear membrane. Other morphological changes that can be measured to determine apoptosis include, *e.g., cytoplasmic condensation, increased membrane blebbing, and cellular shrinkage*.

[0238] The presence of apoptotic cells can be measured in both the attached and "floating"
30 compartments of the cultures. For example, both compartments can be collected by removing the supernatant, trypsinizing the attached cells, combining the preparations following a centrifugation wash step (*e.g., 10 minutes, 2000 rpm*), and detecting apoptosis (*e.g., by measuring DNA fragmentation*). (*See, e.g., Piazza et al., 1995, Cancer Research* 55:3110-16).

35 [0239] Immunosuppressive effects can be measured, for example, by assaying for a cytotoxic or cytostatic effect (as described *supra*) on an activated immune cell involved in promoting

- 5 immune responses (*e.g.*, CD8⁺ cytotoxic T cells, CD4⁺ Th1 cells, B cells, or mature dendritic cells). In addition or alternatively, an immunosuppressive effect can be determined by examination of the presence or absence of specific immune cell populations (using, *e.g.*, flow cytometry) such as, for example, the presence of T regulatory cells involved in suppression of an immune response; measurement of the functional capacity of immune cells, including,
- 10 *e.g.*, the cytolytic capacity of cytotoxic T cells; measurements of the cytokines, chemokines, cell surface molecules, antibodies or other secreted or cell-surface molecules of the cells (*e.g.*, by flow cytometry, enzyme-linked immunosorbent assays, Western blot analysis, protein microarray analysis, immunoprecipitation analysis); or measurement of biochemical markers of activation of immune cells or signaling pathways within immune cells (*e.g.*,
- 15 Western blot and immunoprecipitation analysis of tyrosine, serine or threonine phosphorylation, polypeptide cleavage, and formation or dissociation of protein complexes; protein array analysis; DNA transcriptional profiling using DNA arrays or subtractive hybridization).

20 VI. Animal Models of Immunological Disorders or CD70-Expressing Cancers

[0240] The anti-CD70 ADCs or ADC derivatives can be tested or validated in animal models of immunological disorders or CD70-expressing cancers. A number of established animal models of immunological disorders or CD70-expressing cancers are known to the skilled artisan, any of which can be used to assay the efficacy of the ADC or ADC derivative. Non-

25 limiting examples of such models are described *infra*.

[0241] Some examples for animal models of systemic and organ-specific autoimmune diseases including diabetes, lupus, systemic sclerosis, Sjögren's Syndrome, experimental autoimmune encephalomyelitis (multiple sclerosis), thyroiditis, myasthenia gravis, arthritis, uveitis, inflammatory bowel disease have been described by Bigazzi, "Animal Models of

30 Autoimmunity: Spontaneous and Induced," in *The Autoimmune Diseases* (Rose & Mackay eds., Academic Press, 1998) and in "Animal Models for Autoimmune and Inflammatory Disease," in *Current Protocols in Immunology* (Coligan *et al.* eds., Wiley & Sons, 1997).

[0242] Allergic conditions, *e.g.*, asthma and dermatitis, can also be modeled in rodents. Airway hypersensitivity can be induced in mice by ovalbumin (Tomkinson *et al.*, 2001, *J. Immunol.* 166:5792-5800) or *Schistosoma mansoni* egg antigen (Tesciuba *et al.*, 2001, *J. Immunol.* 167:1996-2003). The Nc/Nga strain of mice show marked increase in serum IgE

5 and spontaneously develop atopic dermatitis-like lesions (Vestergaard *et al.*, 2000, *Mol. Med. Today* 6:209-210; Watanabe *et al.*, 1997, *Int. Immunol.* 9:461-466; Saskawa *et al.*, 2001, *Int. Arch. Allergy Immunol.* 126:239-247).

[0243] Injection of immuno-competent donor lymphocytes into a lethally irradiated histo-incompatible host is a classical approach to induce acute GVHD in mice. Alternatively, the
10 parent B6D2F1 murine model provides a system to induce both acute and chronic GVHD. In this model the B6D2F1 mice are F1 progeny from a cross between the parental strains of C57BL/6 and DBA/2 mice. Transfer of DBA/2 lymphoid cells into non-irradiated B6D2F1 mice causes chronic GVHD, whereas transfer of C57BL/6, C57BL/10 or B10.D2 lymphoid cells causes acute GVHD (Slayback *et al.*, 2000, *Bone Marrow Transpl.* 26:931-938; Kataoka
15 *et al.*, 2001, *Immunology* 103:310-318).

[0244] Additionally, both human hematopoietic stem cells and mature peripheral blood lymphoid cells can be engrafted into SCID mice, and these human lympho-hematopoietic cells remain functional in the SCID mice (McCune *et al.*, 1988, *Science* 241:1632-1639; Kamel-Reid and Dick, 1988, *Science* 242:1706-1709; Mosier *et al.*, 1988, *Nature* 335:256-
20 259). This has provided a small animal model system for the direct testing of potential therapeutic agents on human lymphoid cells. (See, e.g., Tournoy *et al.*, 2001, *J. Immunol.* 166:6982-6991).

[0245] Moreover, small animal models to examine the *in vivo* efficacies of the anti-CD70 ADCs or ADC derivatives can be created by implanting CD70-expressing human tumor cell
25 lines into appropriate immunodeficient rodent strains, e.g., athymic nude mice or SCID mice. Examples of CD70-expressing human lymphoma cell lines include, for example, Daudi (Ghetie *et al.*, 1994, *Blood* 83:1329-36; Ghetie *et al.*, 1990, *Int J Cancer* 15:481-5; de Mont *et al.*, 2001, *Cancer Res* 61:7654-59), Ramos (Ma *et al.*, 2002, *Leukemia* 16:60-6; Press *et al.*, 2001, *Blood* 98:2535-43), HS-Sultan (Cattan & Maung, 1996, *Cancer Chemother*
30 *Pharmacol* 38:548-52; Cattan and Douglas, 1994, *Leuk Res* 18:513-22), Raji (Ochakovskaya *et al.*, 2001, *Clin Cancer Res* 7:1505-10; Breisto *et al.*, 1999, *Cancer Res* 59:2944-49), and CA46 (Kreitman *et al.*, 1999, *Int J Cancer* 81:148-55). Non-limiting example of a CD70-expressing Hodgkin's lymphoma line is L540cy (Barth *et al.*, 2000, *Blood* 95:3909-14; Wahl *et al.*, 2002, *Cancer Res* 62:3736-42). Non-limiting examples of CD70 expressing human
35 renal cell carcinoma cell lines include 786-O (Ananth *et al.*, 1999, *Cancer Res* 59:2210-6; Datta *et al.*, 2001, *Cancer Res* 61:1768-75), ACHN (Hara *et al.*, 2001, *J Urol.* 166:2491-4;

5 Miyake *et al.*, 2002, *J Urol.* 167:2203-8), Caki-1 (Prewett *et al.*, 1998, *Clin. Cancer Res.* 4:2957-66; Shi and Siemann, 2002, *Br. J. Cancer* 87:119-26), and Caki-2 (Zellweger *et al.*, 2001, *Neoplasia* 3:360-7). Non-limiting examples of CD70-expressing nasopharyngeal carcinoma cell lines include C15 and C17 (Busson *et al.*, 1988, *Int. J. Cancer* 42:599-606; Bernheim *et al.*, 1993, *Cancer Genet. Cytogenet.* 66:11-5). Non-limiting examples of CD70-
10 expressing human glioma cell lines include U373 (Palma *et al.*, 2000, *Br. J. Cancer* 82:480-7) and U87MG (Johns *et al.*, 2002, *Int. J. Cancer* 98:398-408). These tumor cell lines can be established in immunodeficient rodent hosts either as solid tumor by subcutaneous injections or as disseminated tumors by intravenous injections. Once established within a host, these tumor models can be applied to evaluate the therapeutic efficacies of the anti-CD70 ADCs or
15 ADC derivatives as described herein on modulating *in vivo* tumor growth.

VII. Immune Disorders and CD70-expressing Cancers

[0246] The anti-CD70 ADCs and ADC derivatives as described herein are useful for treating or preventing an immunological disorder characterized by inappropriate activation of immune
20 cells (*e.g.*, lymphocytes or dendritic cells). Treatment or prevention of the immunological disorder, according to the methods described herein, is achieved by administering to a subject in need of such treatment or prevention an effective amount of the anti-CD70 ADC or ADC derivative, whereby the ADC or ADC derivative (i) binds to activated immune cells that express CD70 and that are associated with the disease state and (ii) exerts a cytotoxic,
25 cytostatic, or immunosuppressive effect on the activated immune cells.

[0247] Immunological diseases that are characterized by inappropriate activation of immune cells and that can be treated or prevented by the methods described herein can be classified, for example, by the type(s) of hypersensitivity reaction(s) that underlie the disorder. These reactions are typically classified into four types: anaphylactic reactions, cytotoxic (cytolytic)
30 reactions, immune complex reactions, or cell-mediated immunity (CMI) reactions (also referred to as delayed-type hypersensitivity (DTH) reactions). (*See, e.g., Fundamental Immunology* (William E. Paul ed., Raven Press, N.Y., 3rd ed. 1993).)

[0248] Specific examples of such immunological diseases include the following: rheumatoid arthritis, autoimmune demyelinating diseases (*e.g.*, multiple sclerosis, allergic
35 encephalomyelitis), endocrine ophthalmopathy, uveoretinitis, systemic lupus erythematosus, myasthenia gravis, Grave's disease, glomerulonephritis, autoimmune hepatological disorder,

- 5 inflammatory bowel disease (*e.g.*, Crohn's disease), anaphylaxis, allergic reaction, Sjogren's syndrome, type I diabetes mellitus, primary biliary cirrhosis, Wegener's granulomatosis, fibromyalgia, polymyositis, dermatomyositis, multiple endocrine failure, Schmidt's syndrome, autoimmune uveitis, Addison's disease, adrenalitis, thyroiditis, Hashimoto's thyroiditis, autoimmune thyroid disease, pernicious anemia, gastric atrophy, chronic hepatitis, lupoid hepatitis, atherosclerosis, subacute cutaneous lupus erythematosus,
- 10 hypoparathyroidism, Dressler's syndrome, autoimmune thrombocytopenia, idiopathic thrombocytopenic purpura, hemolytic anemia, pemphigus vulgaris, pemphigus, dermatitis herpetiformis, alopecia arcata, pemphigoid, scleroderma, progressive systemic sclerosis, CREST syndrome (calcinosis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyly, and telangiectasia), male and female autoimmune infertility, ankylosing spondylitis, ulcerative colitis, mixed connective tissue disease, polyarteritis nodosa, systemic necrotizing vasculitis, atopic dermatitis, atopic rhinitis, Goodpasture's syndrome, Chagas' disease, sarcoidosis, rheumatic fever, asthma, recurrent abortion, anti-phospholipid syndrome, farmer's lung, erythema multiforme, post cardiectomy syndrome, Cushing's
- 20 syndrome, autoimmune chronic active hepatitis, bird-fancier's lung, toxic epidermal necrolysis, Alport's syndrome, alveolitis, allergic alveolitis, fibrosing alveolitis, interstitial lung disease, erythema nodosum, pyoderma gangrenosum, transfusion reaction, Takayasu's arteritis, polymyalgia rheumatica, temporal arteritis, schistosomiasis, giant cell arteritis, ascariasis, aspergillosis, Sampter's syndrome, eczema, lymphomatoid granulomatosis, Behcet's disease, Caplan's syndrome, Kawasaki's disease, dengue, encephalomyelitis, endocarditis, endomyocardial fibrosis, endophthalmitis, erythema elevatum et diutinum, psoriasis, erythroblastosis fetalis, eosinophilic fasciitis, Shulman's syndrome, Felty's syndrome, filariasis, cyclitis, chronic cyclitis, heterochronic cyclitis, Fuch's cyclitis, IgA nephropathy, Henoch-Schonlein purpura, graft versus host disease, transplantation rejection,
- 30 cardiomyopathy, Eaton-Lambert syndrome, relapsing polychondritis, cryoglobulinemia, Waldenstrom's macroglobulemia, Evan's syndrome, and autoimmune gonadal failure.

[0249] Accordingly, the methods described herein encompass treatment of disorders of B lymphocytes (*e.g.*, systemic lupus erythematosus, Goodpasture's syndrome, rheumatoid arthritis, and type I diabetes), Th₁-lymphocytes (*e.g.*, rheumatoid arthritis, multiple sclerosis,

35 psoriasis, Sjogren's syndrome, Hashimoto's thyroiditis, Grave's disease, primary biliary cirrhosis, Wegener's granulomatosis, tuberculosis, or acute graft versus host disease), or Th₂-lymphocytes (*e.g.*, atopic dermatitis, systemic lupus erythematosus, atopic asthma,

5 rhinoconjunctivitis, allergic rhinitis, Omenn's syndrome, systemic sclerosis, or chronic graft versus host disease). Generally, disorders involving dendritic cells involve disorders of Th₁-lymphocytes or Th₂-lymphocytes.

[0250] In certain embodiments, the immunological disorder is a T cell-mediated immunological disorder, such as a T cell disorder in which activated T cell associated with
10 the disorder express CD70. ADC's or ADC derivatives can be administered to deplete such CD70-expressing activated T cells. In a specific embodiment, administration of ADC's or ADC derivatives can deplete CD70-expressing activated T cells, while resting T cells are not substantially depleted by the ADC or ADC derivative. In this context, "not substantially depleted" means that less than about 60%, or less than about 70% or less than about 80% of
15 resting T cells are not depleted.

[0251] The anti-CD70 ADCs and ADC derivatives as described herein are also useful for treating or preventing a CD70-expressing cancer. Treatment or prevention of a CD70-expressing cancer, according to the methods described herein, is achieved by administering to
20 a subject in need of such treatment or prevention an effective amount of the anti-CD70 ADC or ADC derivative, whereby the ADC or ADC derivative (i) binds to CD70-expressing cancer cells and (ii) exerts a cytotoxic or cytostatic effect to deplete or inhibit the proliferation of the CD70-expressing cancer cells.

[0252] CD70-expressing cancers that can be treated or prevented by the methods described herein include, for example, different subtypes of indolent Non-Hodgkin's Lymphoma
25 (indolent NHLs) (*e.g.*, follicular NHLs, small lymphocytic lymphomas, lymphoplasmacytic NHLs, or marginal zone NHLs); cancers of the B-cell lineage, including, *e.g.*, Burkitt's lymphoma and chronic lymphocytic leukemia; renal cell carcinomas; nasopharyngeal carcinomas; thymic carcinomas; gliomas; meningiomas; and colon, stomach, and rectal carcinomas.

30

VIII. Pharmaceutical Compositions Comprising Anti-CD70 ADCs and ADC Derivatives and Administration Thereof

[0253] In accordance with the present methods, a composition comprising an anti-CD70 ADC or ADC derivative as described herein is administered to a subject having or at risk of
35 having an immunological disorder or a CD70-expressing cancer. The term "subject" as used

5 herein means any mammalian patient to which a CD70-binding protein-drug conjugate may be administered, including, *e.g.*, humans and non-human mammals, such as primates, rodents, and dogs. Subjects specifically intended for treatment using the methods described herein include humans. The ADCs or ADC derivatives can be administered either alone or in
10 disorder or CD70-expressing cancer.

[0254] Various delivery systems are known and can be used to administer the anti-CD70 ADC or ADC derivative. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The ADCs or ADC derivatives can be administered, for example by infusion or bolus
15 injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, and the like) and can be administered together with other biologically active agents such as chemotherapeutic agents. Administration can be systemic or local.

[0255] In specific embodiments, the anti-CD70 ADC or ADC derivative composition is administered by injection, by means of a catheter, by means of a suppository, or by means of
20 an implant, the implant being of a porous, non-porous, or gelatinous material, including a membrane, such as a sialastic membrane, or a fiber. Typically, when administering the composition, materials to which the ADC or ADC derivative does not absorb are used.

[0256] In other embodiments, the ADC or ADC derivative is delivered in a controlled release system. In one embodiment, a pump may be used (*see* Langer, 1990, *Science* 249:1527-
25 1533; Sefton, 1989, *CRC Crit. Ref. Biomed. Eng.* 14:201; Buchwald *et al.*, 1980, *Surgery* 88:507; Saudek *et al.*, 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used. (*See Medical Applications of Controlled Release* (Langer & Wise eds., CRC Press, Boca Raton, Florida, 1974); *Controlled Drug Bioavailability, Drug Product Design and Performance* (Smolen & Ball eds., Wiley, New York, 1984); Ranger & Peppas,
30 1983, *Macromol. Sci. Rev. Macromol. Chem.* 23:61. *See also* Levy *et al.*, 1985, *Science* 228:190; During *et al.*, 1989, *Ann. Neurol.* 25:351; Howard *et al.*, 1989, *J. Neurosurg.* 71:105.) Other controlled release systems are discussed, for example, in Langer, *supra*.

[0257] The anti-CD70 ADCs or ADC derivatives are administered as pharmaceutical compositions comprising a therapeutically effective amount of the ADC or ADC derivative
35 and one or more pharmaceutically compatible ingredients. For example, the pharmaceutical composition typically includes one or more pharmaceutical carriers (*e.g.*, sterile liquids, such

5 as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like). Water is a more typical carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include, for example, starch,
10 glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol, and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release
15 formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, *etc.* Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W.
20 Martin. Such compositions will contain a therapeutically effective amount of the nucleic acid or protein, typically in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulations correspond to the mode of administration.

[0258] In typical embodiments, the pharmaceutical composition is formulated in accordance
25 with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the pharmaceutical can also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in
30 unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the pharmaceutical is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the pharmaceutical is administered by injection, an ampoule of sterile water for injection or
35 saline can be provided so that the ingredients can be mixed prior to administration.

[0259] In certain embodiments, the pharmaceutical compositions comprising the anti-CD70 ADC or ADC derivative can further comprise a second therapeutic agent (*e.g.*, a second ADC

5 or ADC derivative or a non-conjugated cytotoxic or immunosuppressive agent such as, for example, any of those described herein).

[0260] Further, the pharmaceutical composition can be provided as a pharmaceutical kit comprising (a) a container containing an anti-CD70 ADC or ADC derivative in lyophilized form and (b) a second container containing a pharmaceutically acceptable diluent (e.g., sterile
10 water) for injection. The pharmaceutically acceptable diluent can be used for reconstitution or dilution of the lyophilized ADC or ADC derivative. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

15 [0261] The amount of the ADC or ADC derivative that is effective in the treatment or prevention of an immunological disorder or CD70-expressing cancer can be determined by standard clinical techniques. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the stage of immunological disorder or CD70-
20 expressing cancer, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0262] For example, toxicity and therapeutic efficacy of the ADCs or ADC derivatives can be determined in cell cultures or experimental animals by standard pharmaceutical
25 procedures for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. ADCs or ADC derivatives that exhibit large therapeutic indices are preferred. Where an ADC or ADC derivative exhibits toxic side effects, a delivery system that targets the ADC or
30 ADC derivative to the site of affected tissue can be used to minimize potential damage non-CD70-expressing cells and, thereby, reduce side effects.

[0263] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of the anti-CD70 ADC or ADC derivative typically lies within a range of circulating concentrations that include the ED₅₀
35 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any ADC or ADC derivative

5 used in the method, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (*i.e.*, the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in
10 plasma can be measured, for example, by high performance liquid chromatography.

[0264] Generally, the dosage of an anti-CD70 ADC or ADC derivative administered to a patient with an immunological disorder or CD70-expressing cancer is typically 0.1 mg/kg to 100 mg/kg of the subject's body weight. More typically, the dosage administered to a subject is 0.1 mg/kg to 50 mg/kg of the subject's body weight, even more typically 1 mg/kg to 30
15 mg/kg, 1 mg/kg to 20 mg/kg, 1 mg/kg to 15 mg/kg, or 1 mg/kg to 10 mg/kg of the subject's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign proteins. Thus, lower dosages of ADCs comprising humanized, chimeric or human antibodies and less frequent administration is often possible.

20 [0265] The anti-CD70 ADC or ADC derivative can be administered in combination with one or more other therapeutic agents for the treatment or prevention of immunological disorders or CD70-expressing cancers. For example, combination therapy can include a second cytostatic, cytotoxic, or immunosuppressive agent (for example, an unconjugated cytostatic, cytotoxic, or immunosuppressive agent such as those conventionally used for the treatment of
25 cancers or immunological disorders). Combination therapy can also include, *e.g.*, administration of an agent that targets a receptor or receptor complex other than CD70 on the surface of activated lymphocytes, dendritic cells or CD70-expressing cancer cells. An example of such an agent is a second, non-CD70 antibody that binds to a molecule at the surface of an activated lymphocyte, dendritic cells or CD70-expressing cancer cells. Another
30 example is a ligand that targets such a receptor or receptor complex. Typically, such an antibody or ligand binds to a cell surface receptor on activated lymphocytes, dendritic cells or CD70-expressing cancer cells and enhances the cytotoxic or cytostatic effect of the anti-CD70 antibody by delivering a cytostatic or cytotoxic signal to the activated lymphocytes, dendritic cells or CD70-expressing cancer cells.

5 [0266] Such combinatorial administration can have an additive or synergistic effect on disease parameters (*e.g.*, severity of a symptom, the number of symptoms, or frequency of relapse).

[0267] With respect to therapeutic regimens for combinatorial administration, in a specific embodiment, an anti-CD70 ADC or ADC derivative is administered concurrently with a
10 second therapeutic agent. In another specific embodiment, the second therapeutic agent is administered prior or subsequent to administration of the anti-CD70 ADC or ADC derivative, by at least an hour and up to several months, for example at least an hour, five hours, 12 hours, a day, a week, a month, or three months, prior or subsequent to administration of the ADC or ADC derivative.

15 [0268] The present invention is not to be limited in scope by the specific embodiments described herein. Various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

20 [0269] The invention is further described in the following examples, which are in not intended to limit the scope of the invention. Cell lines described in the following examples were maintained in culture according to the conditions specified by the American Type Culture Collection (ATCC) or Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany (DMSZ). Cell culture reagents were obtained from
25 Invitrogen Corp., Carlsbad, CA.

Example 1: Sequence Analysis of Anti-CD70 Monoclonal Antibodies 1F6 and 2F2

[0270] To determine the cDNA sequences encoding the light (VL) and heavy (VH) chain variable regions of 1F6 and 2F2 mAb, total RNA was isolated from the 1F6 and 2F2 hybridomas using the TRIzol® Reagent (Invitrogen, Carlsbad, CA) according to the
30 manufacturer's instructions. The gene-specific primers mIgckK1 5'-CTT CCA CTT GAC ATT GAT GTC TTT G-3' (SEQ ID NO:41) and mIgG1 5'-CAG GTC ACT GTC ACT GGC TCA G-3' (SEQ ID NO:42) were applied to reverse transcribe the light chain variable (VL) and heavy chain variable (VH) first strand cDNAs from both RNA preparations, respectively. First strand cDNA reactions were run using the SuperScript™ First Strand Synthesis System
35 for RT-PCR from Invitrogen (Carlsbad, CA). The VL and VH cDNAs were then poly-G tailed using terminal deoxynucleotidyl transferase (TdT) (Invitrogen) and the supplied TdT

5 buffer in conditions specified by the manufacturer. Poly-G tailed V_L and V_H first strand cDNAs were then subjected to PCR amplification. The forward primer for both the V_L V_H PCR was ANCTAIL 5' GTC GAT GAG CTC TAG AAT TCG TGC CCC CCC CCC C-3' (SEQ ID NO:43). The reverse primer for amplifying the V_L was HBS-mck 5'-CGT CAT GTC GAC GGA TCC AAG CTT CAA GAA GCA CAC GAC TGA GGC AC-3' (SEQ ID NO:44).
10 ID NO:44). The reverse primer for amplifying the V_H was HBS-mG1 5'-CGT CAT GTC GAC GGA TCC AAG CTT GTC ACC ATG GAG TTA GTT TGG GC-3' (SEQ ID NO:45). PCR were run with Ex Taq (Fisher Scientific, Pittsburgh, PA) and the supplied reaction buffer in conditions specified by the manufacturer. The V_L and V_H PCR products were then cut by HindIII and EcoRI and cloned into HindIII/EcoRI-cut pUC19. Recombinant plasmid
15 clones were identified, and the nucleotide sequences for the 1F6 and 2F2 hybridomas were determined.


[0271] Complementarity determining regions (CDRs) in the heavy and light chains of 1F6 and 2F2 mAbs were determined according to the criteria described in Kabat *et al.*, 1991, *Sequences of Proteins of Immunological Interest*, Washington DC, US Department of Health
20 and Public Services; Chothia and Lesk, 1987, *J. Mol. Biol.* 196:901-17 (Figures 1 and 2). Sequence alignments at both the cDNA and amino acid levels revealed closely related light chain genes were probably utilized in both hybridomas. There is a 92% sequence identity between 1F6 V_L and 2F2 V_L on the amino acid levels. Sequence comparison of the CDRs shows that 1F6 CDR-L1 is identical to 2F2 CDR-L1, only one divergent substitution is
25 present between 1F6 CDR-L2 and 2F2 CDR-L2, and only 2 conservative substitutions are present between 1F6 CDR-L3 and 2F2 CDR-L3 (Figure 3). On the other hand, a higher degree of sequence diversity is present between 1F6 V_H and 2F2 V_H, about 66 of 137 amino acid residues are different between the 2 V_Hs. Sequence comparison of the CDRs shows that
30 5 of 10 residues are different between 1F6 CDR-H1 and 2F2 CDR-H1 (3 of the 5 substitutions are divergent), 12 of 17 residues are different between 1F6 CDR-H2 and 2F2 CDR-H2 (9 of the 12 substitutions are divergent), and 5 of 9 residues are different between 1F6 CDR-H3 and 2F2 CDR-H3 (4 of the 5 substitutions are divergent) (Figure 3).

Example 2: Synthesis of Anti-CD70 Drug Conjugates

35 [0272] The ability of anti-CD70 to deliver a potent cytotoxic drug in the form of an antibody drug conjugate (ADC) to eliminate CD70-expressing cells was tested. Monomethyl

- 5 auristatin E (MMAE), auristatin phenylalanine phenylenediamine (AFP), and monomethyl auristatin phenylalanine (MMAF) were used as the targeted cytotoxic drugs for this study. The drugs were linked to the anti-CD70 mAb 1F6 by the valine-citrulline (vc) dipeptide linker to give the 1F6-vcMMAE, 1F6-vcAFP conjugates and 1F6-vcMMAF.


A. Synthesis of 1F6-vcMMAE



10

[0273] The synthesis of the ADC 1F6-vcMMAE, the general structure of which is depicted above, is described below.

1. Drug-Linker Compound Synthesis



15

MMAE

- [0274] Synthesis of auristatin E has been previously described (U.S. Patent No. 5,635,483; Pettit, 1999, *Prog. Chem. Org. Nat. Prod.* 70:1-79). The monomethyl derivative of Auristatin E (MMAE) was prepared by replacing a protected form of monomethylvaline for N,N-dimethylvaline in the synthesis of auristatin E (Senter *et al.*, U.S. Provisional Application No. 60/400,403; and PCT/US03/24209 (*supra*)).

- [0275] To prepare the drug-linker compound, MMAE (1.69 g, 2.35 mmol), maleimidocaproyl-L-valine-L-citrulline-*p*-aminobenzyl alcohol *p*-nitrophenylcarbonate (2.6 g, 3.52 mmol, 1.5 eq., prepared as described in Dubowchik *et al.*, 2002, *Bioconjugate Chem.* 13:855-869) and HOBt (64 mg, 0.45 mmol, 0.2 eq.) were diluted with DMF (25 mL). After 25 min, pyridine (5 mL) was added and the reaction was monitored using reverse-phase HPLC. The reaction was shown to be complete in 24 hr. The reaction mixture was concentrated to

5 provide a dark oil, which was diluted with 3 mL of DMF. The DMF solution was purified using flash column chromatography (silica gel, eluant gradient: 100% dichloromethane to 4:1 dichloromethane-MeOH). The relevant fractions were combined and concentrated to provide an oil that solidified under high vacuum to provide a mixture of the desired drug-linker compound and unreacted MMAE as a dirty yellow solid (R_f 0.40 in 9:1 dichloromethane-MeOH).
10 MeOH). The dirty yellow solid was diluted with DMF and purified using reverse-phase preparative-HPLC (Varian Dynamax C18 column 41.4 mm x 25 cm, 8 μ , 100 Å, using a gradient run of MeCN and 0.1% aqueous TFA at 45 mL/min from 10% to 100% over 40 min followed by 100% MeCN for 20 min) to provide the desired drug-linker compound as an amorphous white powder (R_f 0.40 in 9:1 dichloromethane-MeOH) which was >95% pure by
15 HPLC and which contained less than 1% of MMAE. Yield: 1.78 g (57%); ES-MS m/z 1316.7 [M+H]⁺; UV λ_{max} 215, 248 nm.


2. Conjugate Preparation

[0276] *Antibody Reduction.* To 0.7 mL 1F6 (10 mg/mL) was added 90 μ L of 500 mM sodium borate/500 mM NaCl, pH 8.0, followed by 90 μ L of 100 mM DTT in water, and 20
20 μ L of H₂O. After incubation at 37°C for 30 min, the buffer was exchanged by elution over G25 resin equilibrated and eluted with PBS containing 1 mM DTPA (Aldrich). The thiol/Ab value was found to be 9.8 by determining the reduced antibody concentration from the solution's 280 nm absorbance, and the thiol concentration by reaction with DTNB (Aldrich) and determination of the absorbance at 412 nm.

25 [0277] *Conjugation of the Reduced Antibody.* The reduced mAb was chilled on ice. The drug-linker compound was used as a DMSO solution of known concentration, and the quantity of drug-linker added to the reaction mixture was calculated as follows: $L \text{ stock solution} = V \times [Ab] \times \text{Fold Excess} / [\text{Drug-Linker}]$, where V and $[Ab]$ are the volume and molar concentration of the reduced antibody solution, respectively. 0.904 μ L cold H₂O was
30 added to the reduced antibody solution, followed by 500 μ L cold acetonitrile. 30.5 μ L of 10.2 mM drug-linker compound stock solution was diluted into 1.47 mL acetonitrile. The acetonitrile drug-linker solution was chilled on ice, then added to the reduced antibody solution. The reaction was terminated after 0.5 hr by the addition of a 20 fold molar excess of cysteine over maleimide. The reaction mixture was concentrated by centrifugal
35 ultrafiltration and purified by elution through de-salting G25 in PBS at 4°C. 1F6-vcMMAE was then filtered through 0.2 micron filters under sterile conditions and immediately frozen at


- 5 -80°C. 1F6-vcMMAE was analyzed for 1) concentration, by UV absorbance; 2) aggregation, by size exclusion chromatography; 3) drug/Ab, by measuring unreacted thiols with DTNB, and 4) residual free drug, by reverse phase HPLC.

B. Synthesis of 1F6-vcAFP:



- 10 [0278] The synthesis of the ADC 1F6-vcAFP, the general structure of which is depicted above, is described below.

1. AFP Synthesis



AFP

- 15 [0279] Boc-phenylalanine (1.0 g, 3.8 mmol) was added to a suspension of 1,4-diaminobenzene•HCl (3.5 g, 19.0 mmol, 5.0 eq.) in triethylamine (10.7 mL, 76.0 mmol, 20 eq.) and dichloromethane (50 mL). To the resulting solution was added DEPC (3.2 mL, 19.0 mmol, 5.0 eq.) via syringe. HPLC showed no remaining Boc-phe after 24 hr. The reaction mixture was filtered, and the filtrate was concentrated to provide a dark solid. The dark solid residue was partitioned between 1:1 EtOAc-water, and the EtOAc layer was washed sequentially with water and brine. The EtOAc layer was dried and concentrated to provide a dark brown/red residue that was purified using HPLC (Varian Dynamax column 41.4 mm x 25 cm, 5 μ , 100Å using a gradient run of MeCN and water at 45 mL/min from 10% to 100% over 40 min followed by 100% MeCN for 20 min). The relevant fractions were combined and concentrated to provide a red-tan solid intermediate. Yield: 1.4 g (100%); ES-MS m/z 355.9 [M+H]⁺; UV λ max 215, 265 nm; ¹H NMR (CDCl₃) δ 7.48 (1 H, br s), 7.22-7.37 (5 H,
- 20
- 25

5 m), 7.12 (2 H, d, $J=8.7$ Hz), 7.61 (2 H, d, $J=8.7$ Hz), 5.19 (1 H, br s), 4.39-4.48 (1 H, m), 3.49 (2 H, s), 3.13 (2 H, d, $J=5.7$ Hz), 1.43 (9 H, s).

[0280] The red-tan solid intermediate (0.5 g, 1.41 mmol) and diisopropylethylamine (0.37 mL, 2.11 mmol, 1.5 eq.) were diluted with dichloromethane (10 mL), and to the resulting solution was added Fmoc-Cl (0.38 g, 1.41 mmol). The reaction was allowed to stir, and a
10 white solid precipitate formed after a few minutes. Reaction was complete according to HPLC after 1 hr. The reaction mixture was filtered, and the filtrate was concentrated to provide an oil. The oil was precipitated with EtOAc, resulting in a reddish-white intermediate product, which was collected by filtration and dried under vacuum. Yield: 0.75 g (93%); ES-MS m/z 578.1 $[M+H]^+$, 595.6 $[M+NH_4]^+$.

15 [0281] The reddish-white intermediate (0.49 g, 0.85 mmol), was diluted with 10 mL of dichloromethane, and then treated with 5 mL of trifluoroacetic acid. Reaction was complete in 30 min according to reverse-phase HPLC. The reaction mixture was concentrated and the resulting residue was precipitated with ether to provide an off-white solid. The off-white solid was filtered and dried to provide an amorphous powder, which was added to a solution
20 of Boc-Dolaproine (prepared as described in *Tetrahedron*, 1993, 49(9):1913-1924) (0.24 g, 0.85 mmol) in dichloromethane (10 mL). To this solution was added triethylamine (0.36 mL, 2.5 mmol, 3.0 eq.) and PyBrop (0.59 g, 1.3 mmol, 1.5 eq.). The reaction mixture was monitored using reverse-phase HPLC. Upon completion, the reaction mixture was concentrated, and the resulting residue was diluted with EtOAc, and sequentially washed with
25 10% aqueous citric acid, water, saturated aqueous sodium bicarbonate, water, and brine. The EtOAc layer was dried ($MgSO_4$), filtered, and concentrated. The resulting residue was purified using flash column chromatography (silica gel) to provide an off-white powdered intermediate. Yield: 0.57 g (88%); ES-MS m/z 764.7 $[M+NH_4]^+$; UV λ_{max} 215, 265 nm; 1H NMR ($DMSO-d_6$) δ 10.0-10.15 (1 H, m), 9.63 (1 H, br s), 8.42 (1/2 H, d, $J=8.4$ Hz), 8.22 (1/2
30 H, d, $J=8.4$ Hz), 7.89 (2 H, d, $J=7.2$ Hz), 7.73 (2 H, d, $J=7.6$ Hz), 7.11-7.55 (13 H, m), 4.69-4.75 (1 H, m), 4.46 (2 H, d, $J=6.8$ Hz), 4.29 (1 H, t, $J=6.4$ Hz), 3.29 (3 H, s), 2.77-3.47 (7 H, m), 2.48-2.50 (3 H, m), 2.25 (2/3 H, dd, $J=9.6, 7.2$ Hz), 1.41-1.96 (4 H, m), 1.36 (9 H, s), 1.07 (1 H, d, $J=6.4$ Hz, rotational isomer), 1.00 (1 H, d, $J=6.4$ Hz, rotational isomer).


[0282] The white solid intermediate (85 mg, 0.11 mmol) and *N*-Methylval-val-dil-*O*-*t*-butyl
35 (55 mg, 0.11 mmol, prepared as described in Pettit *et al.* 1996, *J. Chem. Soc. Perk. I* p. 859) were diluted with dichloromethane (5 mL), and then treated with 2.5 mL of trifluoroacetic

5 acid under a nitrogen atmosphere for two hours at room temperature. The reaction completion was confirmed by RP-HPLC. The solvent was removed *in vacuo* and the resulting residue was azeotropically dried twice with toluene, and then dried under high vacuum for 12 hours.

[0283] The residue was diluted with dichloromethane (2 mL), diisopropylethylamine (3 eq.) was added, followed by DEPC (1.2 eq.). After the reaction was completed, the reaction mixture was concentrated under reduced pressure, the resulting residue was diluted with EtOAc, and washed sequentially with 10 % aqueous citric acid, water, saturated aqueous sodium bicarbonate, and brine. The EtOAc layer was dried, filtered and concentrated to provide a yellow oil.

15 [0284] The yellow oil was diluted with dichloromethane (10 mL) and to the resulting solution diethylamine (5 mL) was added. According to HPLC, reaction was completed after 2 hr. The reaction mixture was concentrated to provide an oil. The oil was diluted with DMSO, and the DMSO solution was purified using reverse phase preparative-HPLC (Varian Dynamax column 21.4 mm x 25 cm, 5 μ , 100 Å, using a gradient run of MeCN and 0.1% TFA at 20 mL/min from 10% to 100% over 40 min followed by 100% MeCN for 20 min).
20 The relevant fractions were combined and concentrated to provide the desired drug as an off-white solid. Overall yield: 42 mg (44% overall); ES-MS m/z 837.8 $[M+H]^+$, 858.5 $[M+Na]^+$; UV λ_{max} 215, 248 nm.

2. Preparation of Drug-Linker Compound



[0285] The trifluoroacetate salt of AFP (0.37 g, 0.39 mmol, 1.0 eq.) and Fmoc-val-cit (0.30 g, 0.58 mmol, 1.5 eq., prepared according to Dubowchik *et al.*, 2002, *Bioconjugate Chem.* 13:855-896) were diluted with DMF (5 mL, 0.1 M), and to the resulting solution was added pyridine (95 μ L, 1.2 mmol, 3.0 eq.). HATU (0.23 g, 0.58 mmol, 1.5 eq.) was then added as a solid, and the reaction mixture was allowed to stir under argon atmosphere while being monitored using HPLC. The reaction progressed slowly, and 4 hr later, 1.0 eq. of diisopropylethylamine was added. Reaction was complete in 1 hr. The reaction mixture was

5 concentrated *in vacuo* and the resulting residue was purified using prep-HPLC (Varian Dynamax C18 column 41.4 mm x 25 cm, 5 μ , 100 Å, using a gradient run of MeCN and 0.1% aqueous TFA at 45 mL/min from 10% to 100% over 40 min followed by 100% MeCN for 20 min) to provide a faint pink solid intermediate.

[0286] The pink solid intermediate was diluted with DMF (30 mL) and to the resulting
10 solution was added diethylamine (15 mL). Reaction was complete by HPLC in 2 hr. The reaction mixture was concentrated and the resulting residue was washed twice with ether. The solid intermediate was dried under high vacuum and then used directly in the next step.

[0287] The solid intermediate was diluted with DMF (20 mL) and to the resulting solution was added 6-(2,5-dioxy-2,5-dihydro-pyrrol-1-yl)-hexanoic acid 2,5-dioxy-pyrrolidin-1-yl
15 ester (0.12 g, 0.39 mmol, 1.0 eq.) (EMCS, Molecular Biosciences Inc., Boulder, CO). After 4 days, the reaction mixture was concentrated to provide an oil which was purified using prep-HPLC (Varian Dynamax C18 column 41.4 mm x 25 cm, 5 μ , 100 Å, using a gradient run of MeCN and 0.1% aqueous TFA at 45 mL/min from 10% to 100% over 40 min followed by 100% MeCN for 20 min) to provide the desired drug-linker compound as a white flaky solid.
20 Yield: 0.21 g (38% overall); ES-MS m/z 1285.9 $[M+H]^+$; 13.07.8 $[M+Na]^+$; UV λ_{max} 215, 266 nm.


3. Conjugate Preparation

[0288] *Antibody Reduction.* To 0.48 mL 1F6 (10.4 mg/mL) was added 75 μ L of 500 mM sodium borate/500mM NaCl, pH 8.0, followed by 75 μ L of 100 mM DTT in water, and 20
25 μ L PBS. After incubation at 37°C for 30 min, the buffer was exchanged by elution over G25 resin equilibrated and eluted with PBS containing 1 mM DTPA (Aldrich). The thiol/Ab value was found to be 10.3 by determining the reduced antibody concentration from the solution's 280 nm absorbance, and the thiol concentration by reaction with DTNB (Aldrich) and determination of the absorbance at 412 nm.

30 [0289] *Conjugation of the Reduced Antibody.* The reduced mAb was chilled on ice. The drug-linker compound was used as a DMSO solution of known concentration, and the quantity of drug-linker added to the reaction mixture was calculated as follows: L stock solution = $V \times [Ab] \times \text{Fold Excess}/[\text{Drug-Linker}]$, where V and [Ab] are the volume and molar concentration of the reduced antibody solution, respectively. 984 μ L cold PBS/DTPA
35 was added to the reduced antibody solution, followed by 400 μ L acetonitrile, and the mixture chilled on ice. 26.4 μ L of 8.3 mM drug-linker compound stock solution was then added to


- 5 the reduced antibody/DMSO solution. The reaction was terminated after 1 hr by the addition of a 40 μ L of 100 mM cysteine. The reaction mixture was concentrated by centrifugal ultrafiltration and purified by elution through de-salting G25 in PBS at 4°C. 1F6-vcAFP was then filtered through 0.2 micron filters under sterile conditions and immediately frozen at -80°C. 1F6-vcAFP was analyzed for 1) concentration, by UV absorbance; 2) aggregation, by size exclusion chromatography; 3) drug/Ab, by measuring unreacted thiols by treatment with DTT, followed by DTNB, and 4) residual free drug, by reverse phase HPLC.

C. Synthesis of 1F6-vcMMAF



- [0290] The synthesis of the ADC 1F6-vcMMAF, the general structure of which is depicted above, is described below.

1. MMAF Synthesis




Compound 1

- [0291] MMAF was prepared from its *t*-butyl ester (Compound 1) as is described below. Phenylalanine *t*-butyl ester HCl salt (868 mg, 3 mmol), *N*-Boc-Dolaproine (668 mg, 1 eq.), DEPC (820 μ L, 1.5 eq.), and DIEA (1.2 mL) were diluted with dichloromethane (3 mL). After 2 hours (h) at room temperature (about 28 degrees Celcius), the reaction mixture was diluted with dichloromethane (20 mL), washed successively with saturated aqueous (aq.) NaHCO₃ (2 x 10 mL), saturated aq. NaCl (2 x 10 mL). The organic layer was separated and concentrated. The resulting residue was re-suspended in ethyl acetate and was purified via flash chromatography on silica gel in ethyl acetate. The relevant fractions were combined and concentrated to provide the dipeptide as a white solid: 684 mg (46 %). ES-MS m/z 491.3 [M+H]⁺.

5 [0292] For selective Boc cleavage in the presence of *t*-butyl ester, the above dipeptide (500 mg, 1.28 mmol) was diluted with dioxane (2 mL). 4M HCl/dioxane (960 μ L, 3 eq.) was added, and the reaction mixture was stirred overnight at room temperature. Almost complete Boc deprotection was observed by RP-HPLC with minimal amount of *t*-butyl ester cleavage. The mixture was cooled down on an ice bath, and triethylamine (500 μ L) was added. After 10
10 min., the mixture was removed from the cooling bath, diluted with dichloromethane (20 mL), washed successively with saturated aq. NaHCO₃ (2 x 10 mL), saturated aq. NaCl (2 x 10 mL). The organic layer was concentrated to give a yellow foam: 287 mg (57 %). The intermediate was used without further purification.

[0293] The tripeptide *N*-Fmoc-*N*-Methylval-val-dil-*O*-*t*-butyl (0.73 mmol) was treated
15 with TFA (3 mL), dichloromethane (3 mL) for 2 h at room temperature. The mixture was concentrated to dryness, the residue was co-evaporated with toluene (3 x 20 mL), and dried in vacuum overnight. The residue was diluted with dichloromethane (5 mL) and added to the deprotected dipeptide Dap-phe-*O*-*t*-butyl (287 mg, 0.73 mmol), followed by DIEA (550 μ L, 4 eq.), DEPC (201 μ L, 1.1 eq.). After 2 h at room temperature the reaction mixture was
20 diluted with ethyl acetate (50 mL), washed successively with 10% aq. citric acid (2 x 20 mL), saturated aq. NaHCO₃ (2 x 10 mL), saturated aq. NaCl (10 mL). The organic layer was separated and concentrated. The resulting residue was re-suspended in ethyl acetate and was purified via flash chromatography in ethyl acetate. The relevant fractions were combined and concentrated to provide *N*-Fmoc-*N*-Methylval-val-dil-dap-phe-*O*-*t*-butyl as a white solid: 533
25 mg (71 %). R_f 0.4 (EtOAc). ES-MS *m/z* 1010.6 [M+H]⁺.

[0294] The product (200 mg, 0.2 mmol) was diluted with dichloromethane (3 mL), diethylamine (1 mL). The reaction mixture was stirred overnight at room temperature. Solvents were removed to provide an oil that was purified by flash silica gel chromatography in a step gradient 0-10 % MeOH in dichloromethane to provide Compound 1 as a white
30 solid: 137 mg (87 %). R_f 0.3 (10 % MeOH/CH₂Cl₂). ES-MS *m/z* 788.6 [M+H]⁺.



- 5 [0295] MMAF was prepared from Compound 1 (30 mg, 0.038 mmol) by treatment with 4M HCl/dioxane (4 ml) for 7 h at room temperature. The solvent was removed, and the residue was dried in a vacuum overnight to give provide MMAF as a hygroscopic white solid: 35 mg (120 % calculated for HCl salt). ES-MS m/z 732.56 $[M+H]^+$.

2. Preparation of Drug-Linker Compound

- 10 [0296] Compound 1 (83 mg, 0.11 mmol), maleimidocaproyl-L-valine-L-citrulline-*p*-aminobenzyl alcohol *p*-nitrophenylcarbonate (85 mg, 0.12 mmol, 1.1 eq.), and HOBt (2.8 mg, 21 μ mol, 0.2 eq.) were taken up in dry DMF (1.5 mL) and pyridine (0.3 mL) while under argon. After 30 h, the reaction was found to be essentially complete by HPLC. The mixture was evaporated, taken up in a minimal amount of DMSO and purified by prep-HPLC (C₁₂-
15 RP column, 5 μ , 100 Å, linear gradient of MeCN in water (containing 0.1% TFA) 10 to 100 % in 40 min followed by 20 min at 100 %, at a flow rate of 25 mL/min) to provide MC-Val-Cit-PAB-MMAF-O-*t*-butyl ester as a white solid. Yield: 103 mg (71%). ES-MS m/z 1387.06 $[M+H]^+$, 1409.04 $[M+Na]^+$; UV λ_{max} 205, 248 nm.

- [0297] MC-Val-Cit-PAB-MMAF-O-*t*-butyl ester (45 mg, 32 μ mol) was suspended in
20 methylene chloride (6 mL) followed by the addition of TFA (3 mL). The resulting solution stood for 2 h. The reaction mixture was concentrated *in vacuo* and purified by prep-HPLC (C₁₂-RP column, 5 μ , 100 Å, linear gradient of MeCN in water (containing 0.1% TFA) 10 to 100 % in 40 min followed by 20 min at 100 %, at a flow rate of 25 mL/min). The desired fractions were concentrated to provide MC-Val-Cit-PAB-MMAF as an off-white solid.
25 Yield: 11 mg (25%). ES-MS m/z 1330.29 $[M+H]^+$, 1352.24 $[M+Na]^+$; UV λ_{max} 205, 248 nm.

3. Conjugate Preparation

- [0298] *Antibody Reduction.* To 0.75 mL 1F6 (4.2 mg/mL) was added 50 μ L of 500 mM
30 sodium borate/500mM NaCl, pH 8.0, followed by 100 μ L of 100 mM DTT in water, and 100 μ L PBS. After incubation at 37°C for 30 min, the buffer was exchanged by elution over G25 resin equilibrated and eluted with PBS containing 1 mM DTPA (Aldrich). The thiol/Ab value was found to be 8.9 by determining the reduced antibody concentration from the solution's 280 nm absorbance, and the thiol concentration by reaction with DTNB (Aldrich)
35 and determination of the absorbance at 412 nm.

5 [0299] *Conjugation of the Reduced Antibody.* The reduced mAb was chilled on ice. The drug-linker compound was used as a DMSO solution of known concentration, and the quantity of drug-linker added to the reaction mixture was calculated as follows: $L \text{ stock solution} = V \times [Ab] \times \text{Fold Excess} / [\text{Drug-Linker}]$, where V and [Ab] are the volume and molar concentration of the reduced antibody solution, respectively. The reduced antibody
10 (785 μl of 1.77 mg/ml) was added to the mixture of 7.2 μl of 12.9 mM MCvcMMAF in 92 μl of acetonitrile and mixed rapidly. The reaction mixture was allowed to incubate on ice for 1h and followed by addition of a 40 μL of 100 mM cysteine. The reaction mixture was purified by G25 column equilibrated in PBS at 4°C. 1F6-vcAFP was then filtered through 0.2 micron filters under sterile conditions and immediately frozen at -80°C. 1F6-vcMMAF was analyzed
15 for 1) concentration, by UV absorbance; 2) aggregation, by size exclusion chromatography; 3) drug/Ab, by measuring unreacted thiols by treatment with DTT, followed by DTNB, and 4) residual free drug, by reverse phase HPLC.

Example 3: Expression of CD70 on Hematologic Cell Lines

[0300] Surface expression of CD70 was examined using flow cytometry. In general, 0.2×10^6 cells were incubated with 50 μl of staining medium (RPMI-1640 supplemented with 5-
20 10% FBS) containing a fluorochrome-conjugated mAb (10 $\mu\text{g/ml}$) for single-color flow cytometry or a cocktail of fluorochrome-conjugated mAbs for multiple-color flow cytometry. Incubations were carried out on ice for 20-30 minutes. Cells were then washed 3 times with the staining buffer and fixed in PBS containing 1% of paraformaldehyde. Flow cytometric
25 analysis was performed with a FACScan (BD Immunocytometry, San Jose, CA) and data were analyzed by either the CellQuest (BD Immunocytometry) or the WinMDI software. The anti-human CD70 monoclonal antibodies (mAb) 1F6 and 2F2 were obtained from the Central Laboratory of the Netherlands Red Cross Blood Transfusion Service (Amsterdam, The Netherlands). 1F6 was conjugated to AlexaFluor488 (AF) (Molecular Probes, Eugene,
30 OR) according to the manufacturer's instruction, and AF-conjugated 1F6 was used to detect CD70 by flow cytometry whenever necessary. The anti-CD70 mAb Ki-24 and anti-CD3 mAb were purchased from BD PharMingen (San Diego, CA).

[0301] The expression of cell surface CD70 was surveyed on a panel of hematologic cell lines. This panel included both Epstein-Barr Virus (EBV) negative American Burkitt's
35 lymphoma (BL) and EBV⁺ African Burkitt's lymphoma lines, a variety of non-Hodgkin's lymphoma (NHL) lines which are EBV⁺, 2 Hodgkin's disease (HD) lines, 2 EBV-transformed

5 B lymphoblastoid cell lines (EBV-LCL), and an acute T cell leukemia line Jurkat (Figure 4). The cell lines were maintained in culture conditions specified by the ATCC or DSMZ. For flow cytometric analysis, AF-conjugated or a commercially available anti-CD70 (clone Ki-24) purchased from BD PharMingen was used. Cell lines on which anti-CD70 showed greater than 2-fold binding compared to the control IgG binding, based on mean fluorescence intensities, were arbitrarily defined as CD70-expressing. Based on this criterion, all of the BL lines, both EBV⁺ and EBV⁻, examined expressed detectable, albeit low levels of CD70. Five out ten NHL cell lines expressed CD70. Both HD and EBV-LCL also expressed CD70 at levels that were generally higher than those of BL and NHL lines. The acute T leukemia cell line Jurkat was found to be CD70⁺.

15
Example 4: Proliferation Inhibitory and Cytotoxic Effects of Anti-CD70 ADCs on CD70⁺ Hematologic Lines

[0302] A subset of the CD70⁺ hematologic lines listed in Figure 4 was tested for sensitivity to the 1F6 ADCs. These lines were seeded at 5,000 cells per well in 96-well plates in quadruplicates in a total of 200 µl of culture medium containing graded doses of the ADCs as indicated in the Figure 5. Proliferation assays were carried out for 96 hours. Tritiated thymidine (³H-TdR) incorporation during the last 16 hours of incubation was used to assess DNA synthesis. The responses of a CD70⁺ B lymphoma cell line WSU-NHL and the CD70⁺ Jurkat cells are shown in the upper panels of Figure 5. Both 1F6-vcMMAE and 1F6-vcAFP exerted dose-dependent inhibitory effects on the proliferation of WSU-NHL cells. Proliferation inhibitory effects were apparent when the ADCs were used at concentrations higher than 0.005 µg/ml. The non-binding IgG control ADCs did not exert any significant effects at concentrations lower than 0.5 µg/ml. On the other hand, proliferation of the CD70⁺ Jurkat acute T leukemia cell line was not affected by either 1F6-vcMMAE or 1F6-vcAFP at concentrations as high as 2 µg/ml. Hence, the proliferation inhibitory activity of the 1F6 ADCs on target positive cells resulted from specific binding of ADCs to CD70 expressed on the target cells. The responses of other CD70⁺ cell lines tested are summarized in the lower panel of Figure 5. Significant proliferation inhibition by the 1F6 ADCs was observed in 2 of the 3 B lineage non-Hodgkin's lymphoma cell lines (MC116 and WSU-NHL), 2 of 2 Hodgkin's disease lines, and the EBV-transformed B lymphoblastoid cell line CESS. In all the responsive cell lines tested, 1F6-vcAFP was demonstrated to exert more potent inhibitory

5 activity than 1F6-vcMMAE. These results suggest that anti-CD70 ADCs can potentially be applied in immunotherapy of CD70⁺ lymphomas. Moreover, EBV-LCLs are phenotypically very similar to activated normal B cells. Both EBV-LCLs and activated normal B cells are also efficient antigen presenting cells possessing strong T cell stimulatory activities. The observation that anti-CD70 ADC can efficiently suppress the proliferation of EBV-LCL
10 CESS cells indicates that anti-CD70 ADCs can also be applied to suppress immune responses by eliminating activated B cells.

Example 5: Expression of CD70 Transcripts in Solid Tumors

[0303] Hodgkin's disease and nasopharyngeal carcinoma (NPC) are two examples of
15 conditions that frequently associate with EBV and a prominent lymphoid stroma at the tumor sites. The role of this lymphoid stroma is uncertain and may represent infiltration of lymphocytes to tumors as part of the immune response to tumor-associated antigens. Alternatively, the recruitment of lymphocytes by tumor cells may be a mechanism through which tumor cells derive cytokine and growth factors from the infiltrating lymphocytes in a
20 paracrine fashion. Some of these growth factors may contribute to tumor development by supporting tumor cell proliferation as hypothesized for HD (Gruss *et al.*, 1997, *Immunol. Today* 18:156-63). CD70 has been found to be present on the Reed-Sternberg cells in HD, and Figure 6 demonstrates the expression of CD70 on HD cell lines. In a study on frozen tumor biopsies from EBV-associated undifferentiated nasopharyngeal carcinomas, 80% (16
25 of 20 cases) showed *in situ* expression of the CD70 protein on tumor cells (Agathangelou *et al.*, 1995, *Am. J. Pathol.* 147:1152-60). Others also reported the expression of CD70 on EBV- tumors including thymic carcinomas (Hishima *et al.*, 2000, *Am. J. Surg. Path.* 24:742-746), gliomas, and meningiomas (Held-Feindt and Mentlein, 2002, *Int. J. Cancer* 98:352-56).

[0304] The potential expression of CD70 in additional carcinoma types was further
30 surveyed using the Cancer Profiling Array (CPA) (BD Biosciences Clontech, Palo Alto, CA). The CPA includes normalized cDNA from 241 tumor and corresponding normal tissues from individual patients. In order to examine the expression of CD70 message in these patients, a piece of cDNA corresponding to 3' untranslated region of nucleotides 734-874 of the CD70 message was amplified using the reverse transcriptase polymerase chain reaction (RT-PCR)
35 approach. First strand cDNA was synthesized from total RNA isolated from the Burkitt's lymphoma cell line Ramos (ATCC) using the SuperScript™ First Strand Synthesis System

5 for RT-PCR from Invitrogen (Carlsbad, CA). The forward primer 5'-CCA CTG CTG CTG
ATT AG-3' (SEQ ID NO:46), the reverse primer 5'-CAA TGC CTT CTC TTG TCC-3'
(SEQ ID NO:47), and the Advantage 2 PCR Kit (Clontech, Palo Alto, CA) were used for the
PCR. The PCR product was cloned into the pCR4-TOPO vector and sequence verified. To
10 generate a probe for hybridization, PCR was run using the cloned cDNA as the template, the
above primer pairs, and the Advantage 2 PCR Kit (Clontech). To label the probe with ³²P-
dGTP, 200 ng of the purified PCR product were combined with 75 ng of random hexamer, 33
μM each of dATP, dTTP and dCTP, 5 μl of α-³²P dGTP (approximately 3000 Ci/mmol, 10
mCi/ml, Amersham Pharmacia Biotech, Piscataway, NJ), 1 μl of Klenow fragment (New
England Biolabs, Beverly, MA), and 1X EcoPol buffer (New England Biolabs) in a total
15 volume of 50 μl. The reaction mixture was incubated at room temperature for 15 minutes,
and EDTA was added to a final concentration of 10 mM to stop the reaction. Labeled probe
was purified using ProbeQuant G-50 Micro Columns (Amersham Pharmacia Biotech,
Piscataway, NJ). This probe was hybridized to patient cDNA spotted on a CPA using the BD
ExpressHyb hybridization solution (BD Biosciences) according to the manufacturer's
20 instructions. Hybridization signals were quantified by phospho-imaging on a PhosphorImager
SI (Amersham). Reprobing the same CPA for the housekeeping EF-1 gene gave
tumor:normal ratio of approximately one for each sample pair, confirming comparable
loading of cDNA. Therefore ratios between the CD70 hybridization signals obtained from
tumor and normal cDNAs were used as a semi-quantitative measurement to CD70 transcript
25 expression.

[0305] Differential hybridization of the CD70 probe to tumor cDNA was observed in
several cancer types. Most notably, cDNAs from 9 of 20 cases of kidney carcinoma and one
case each of colon, stomach, and rectum carcinoma showed more intense hybridization
signals than cDNAs from the corresponding normal tissues. The 9 cases (45%) of kidney
30 cancer in Figure 6, all classified as RCC, showed more than 2-fold (range of 2.3 to 8.9) over-
expression of CD70 transcripts. In the single case of stomach, colon, and rectum cancer
showing differential CD70 cDNA hybridization (Figure 6), the tumor:normal ratios were 2.9,
8.9, and 3.2, respectively.

[0306] Quantitative PCR (QPCR) was used to quantify CD70 transcript expression in 8
35 RCC and 4 normal kidney cDNA samples, all obtained from BioChain (Hayward, CA). The
RCC cDNA samples were from donors independent of those on the CPA in Figure 6. QPCR
analysis was performed using TaqMan™ MGB chemistry (Applied Biosystems, Foster City,

5 CA) containing a reporter dye at the 5' end (FAM or VIC) and a non-fluorescent quencher (NFQ) at the 3' end on an ABI PRISM 7000 Sequence Detection System (Applied Biosystems). Assays specific for either CD70 (Applied Biosystems Assay# Hs00174297_m1) or control gene GAPDH (Applied Biosystems PDAR# 4326317E) were used to evaluate samples. Comparison of CD70 transcript expression between normal kidney and RCC tissues was conducted based on the comparative threshold cycle (Ct) method as previously reported (Winer *et al.*, 1999, *Anal. Biochem.* 270: 41-49; Aarskog and Vedeler, 2000, *Hum. Genet.* 107: 494-498). Ct values for the CD70 and a housekeeping gene GAPDH from two independent reactions were generated. The Ct values for the GAPDH transcripts from normal kidney and RCC samples did not differ significantly from each other (<1 cycle, student's t-test: $p=0.405$), suggesting that CD70 expression can be normalized against GAPDH for tumor:normal comparison. ΔCt values for CD70 expression in RCC and normal kidney samples were calculated by subtracting Ct_{GAPDH} from Ct_{CD70} to give ΔCt_{Tumor} and ΔCt_{Normal} , respectively. $\Delta\Delta Ct$ values were calculated by subtracting ΔCt_{Normal} from ΔCt_{Tumor} . CD70 transcript expression in RCC expressed as fold increase over normal kidneys was then expressed as $2^{(-\Delta\Delta Ct)}$. CD70 transcript over-expression (>2-fold) was observed in 6 of the 8 RCC cDNA samples, consistent with the results obtained from the CPA experiment (Figure 6). Over-expression of CD70 transcripts ranged from 2.5- to 133.7-fold (Figure 7).

Example 6: Expression of CD70 Protein on Renal Cell Carcinoma

25 [0307] Immunohistochemistry was used to determine if over-expression of CD70 transcript in RCC was paralleled by CD70 protein expression. Frozen RCC tissue sections and normal adjacent tissue sections from two patients independent of those used in CD70 transcript analysis (Figures 6 and 7) were fixed in acetone at -20°C for 10 minutes. After rehydration in PBS, sections were blocked with PBS containing 5% normal goat serum and Avidin D for 30 minutes at room temperature, and washed in PBS. Sections were incubated with the anti-CD70 mAb 2F2 or non-binding IgG at 2 $\mu g/ml$ in PBS containing 5% normal goat serum with biotin for 60 minutes at room temperature. After washes with PBS, biotinylated anti-mouse IgG (VECTASTAIN ABC Kit, Vector Laboratories, Burlingame, CA) at a 1:250 dilution in PBS containing 5% normal goat serum was used to detect bound primary antibody by incubating for 30 minutes at room temperature. Excess biotinylated anti-mouse IgG was removed by PBS washes. Sections were quenched with 1% H_2O_2 in PBS for 30 minutes at room temperature, washed, and then incubated with the VECTASTAIN ABC complex made

5 up according to the manufacturer's instructions. After PBS washes, sections were incubated for 30 minutes at room temperature with the DAB solution (VECTASTAIN ABC Kit) made up according to the manufacturer's instructions for color development. Sections were then washed in water, counterstained with hematoxylin for 2 minutes, mounted in VectaMount (Vector Laboratories), and then observed using 40X light microscopy.

10 [0308] Intense binding of 2F2 to RCC tumor sections from 2 different donors was observed (Figure 8A, right panels). In contrast, a control IgG did not bind to serial tumor sections, confirming the specificity of 2F2 for CD70 (Figure 8A, left panels). 2F2 also did not demonstrate binding beyond that of the control IgG to normal adjacent tissues from the same RCC samples (Figure 8B), suggesting minimal or no CD70 protein expression in normal
15 kidneys. These results confirm the over-expression of CD70 transcripts in RCC and provide direct evidence for *in situ* CD70 protein expression in RCC.

[0309] Expression of the CD70 protein RCC cell surface was next examined. A panel of 7 RCC lines was evaluated using flow cytometry. The RCC lines Hs 835.T, Caki-1, Caki-2, 786-O, 769-P, and ACHN were obtained from the ATCC (Manassas, VA) and were
20 maintained in conditions specified by the ATCC. The RCC lines CAL54 and A498 were obtained from Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (Braunschweig, Germany). Cell lines were maintained in the conditions specified the vendors. The RCC lines SK-RC-6 and SK-RC-7 have been reported previously (Murakami *et al.*, 1984, *Hepatology* 42:192-8). SK-RC-6 and SK-RC-7 was maintained in DMEM
25 supplemented with 10% fetal bovine serum (FBS). Two normal human renal tubule epithelial lines, RPTEC and HRCE (Cambrex, East Rutherford, NJ) were also included in the analysis. Figure 9A shows that the RCC lines 786-O, Caki-1, and Caki-2 expressed CD70 while Hs 835.T did not. Low, but detectable, levels of CD70 expression were also observed on the normal kidney tubule epithelial lines RPTEC and HRCE. Figure 9B summarizes the
30 relative levels of CD70 expressed on RCC lines and normal kidney epithelial cells lines based on the mean fluorescence intensity obtain from flow cytometry. Of the 10 RCC lines tested 9 were found to be CD70 expressing. These results demonstrate that the presence of CD70 transcripts and protein in kidney carcinomas was accompanied by surface expression of CD70 on the cancer cells.

35

5 Example 7: Proliferation Inhibitory and Cytotoxic Effects of Anti-CD70 ADCs on CD70⁺
Renal Cell Carcinoma Cell Lines

[0310] To examine the responses of CD70⁺ RCC lines to 1F6 ADCs, cells were seeded between 1,000 to 3,000 cells per well in 100 μ l of medium. An additional 100 μ l of culture medium containing graded doses of ADCs were added to the wells after cells were allowed to
10 attached overnight. In addition to DNA synthesis, cell viability, as a reflection of the cytotoxic activities of ADCs, was also assayed by the reduction of alamarBlue™ (Biosource International, Camarillo, CA). The alamarBlue™ dye was added at a one to four dilution during the last 4 to 24 hours of the incubation. Dye reduction was assessed by fluorescence spectrometry using the excitation and emission wavelengths of 535 nm and 590 nm,
15 respectively. For analysis, the amount of ³H-TdR incorporated or the extent of alamarBlue™ reduction by the treated cells was compared to that of the untreated control cells. Figure 10 shows the effects of 1F6-vcAFP on the proliferation (³H-TdR incorporation) and viability (alamarBlue™ reduction) on two representative CD70⁺ RCC lines. 1F6-vcAFP demonstrated potent proliferation inhibitory (Figure 10, upper panel) and cytotoxic (Figure 10, lower panel)
20 activities on Caki-1 and 786-O cells, maximal effects were achieved at concentrations lower than 0.1 μ g/ml. In contrast, non-binding control IgG-vcAFP at concentrations lower than 1 μ g/ml exerted minimal effects on these cells.

[0311] Figure 11 summarizes the IC₅₀s for 1F6-vcMMAE, 1F6-vcAFP, and 1F6-vcMMAF on inhibiting the proliferation and inducing cytotoxicity in RCC and normal kidney tubule
25 epithelial cells. 1F6-vcMMAE was very effective in suppressing the proliferation of Caki-1 and Cak-2 cells with IC₅₀ values lower than 20 ng/ml. However, it was not active on other CD70⁺ RCC lines, the CD70⁻ Hs 835.T cells, or the normal kidney epithelial lines RPTEC and HRCE. IC₅₀ values for proliferation inhibition by 1F6-vcAFP ranged from 2-247 ng/ml for CD70⁺ RCC lines. Substantially higher IC₅₀ values were obtained for 1F6-vcAFP on the
30 CD70⁻ Hs 835.T, suggesting specific targeting of CD70. 1F6-vcMMAF were also active in inhibiting proliferation in all CD70⁺ RCC lines tested, with IC₅₀ values lower than 30 ng/ml. Proliferation inhibition was paralleled with cytotoxicity. IC₅₀ values for 1F6-vcAFP to induce cytotoxicity were below 20 ng/ml in 4 of the 7 CD70⁺ RCC lines tested, while those for 1F6-vcMMAF were below 30 ng/ml in 5 of 7 CD70⁺ RCC lines tested. Specific targeting by the
35 1F6 ADC to CD70 was confirmed by the much higher IC₅₀ values obtained for the control non-binding IgG ADCs, >1000 ng/ml for all the lines tested. Moreover, RCC appeared to be differentially more sensitive to anti-CD70 ADC-mediated proliferation inhibition and

- 5 cytotoxicity, as despite detectable expression of CD70, normal kidney epithelial cells were very insensitive to 1F6 ADCs when compared to the CD70⁺ RCC lines.

Example 8: *In Vivo* Efficacy of 1F6-vcAFP in a Xenograft Model of RCC

- [0312] Subcutaneous xenografts of Caki-1 cells have been successfully used as models for
10 RCC to test the efficacy of antibody-based therapeutics including an anti-VEGF antibody (Dagnaes-Hansen *et al.*, 2003, *Anticancer Res.* 23: 1625-30) as well as an ADC consisting of calicheamicin thetal1 conjugated to an anti-gamma-glutamyltransferase antibody (Knoll *et al.*, 2000, *Cancer Res.* 60: 6089-94). Using xenografts of the Caki-1 line in nude mice, the *in vitro* antitumor activity of 1F6-vcAFP was tested. To establish Caki-1 tumors, nude mice
15 were injected subcutaneously with 5×10^6 Caki-1 cells in 0.2 ml PBS. Average tumor size increased to more than 500 mm^3 within 55 days post injection. Caki-1-containing tumors were excised from tumor bearing mice and tumor tissue blocks of approximately 30 mm^3 were prepared. Naïve nude mice to be used for evaluating *in vivo* antitumor activity of anti-CD70 ADC were each implanted subcutaneously with one tumor block of 30 mm^3 .
20 Following implantation, tumor sizes increased to more than 800 mm^3 within 40 days in the absence of any treatment (Figure 12). Treatment was initiated when the average tumor size within a group was approximately 100 mm^3 (Figure 12, arrows). Very little therapeutic activity was detected in 1F6, as tumor growth rate in mice treated with 1F6 was virtually identical to either the untreated control or the IgG-treated group. However, treatment of mice
25 with 1F6-vcAFP significantly inhibited tumor growth. The average tumor size in mice treated with 1 mg/ml of 1F6-vcAFP was $<600 \text{ mm}^3$ 56 days after treatment initiation, whereas in the group treated with IgG-vcAFP at the same dose the average tumor size was already $>600 \text{ mm}^3$ 20 days after treatment initiation (Figure 10, upper panel). Increasing the dose of 1F6-vcAFP to 3 mg/ml further suppressed tumor growth; average tumor size was
30 kept under 200 mm^3 70 days after treatment initiation while IgG-vc-AFP at the same dose resulted in tumors $>400 \text{ mm}^3$ 42 days after treatment initiation (Figure 12, upper panel). The potent, specific *in vivo* antitumor activity of 1F6-vcAFP at 3 mg/kg was from a separate experiment is shown in Figure 12, lower panel.

5 Example 9: Expression of CD70 on Activated T Cells

[0313] The generation and maintenance of the T cell clones have been described in U.S. Provisional Patent Application No. 60/331,750 and in International Patent Application PCT/US02/37223, both of which are incorporated herein by reference in their entireties. To activate resting T cell clones, 5×10^6 T cells were incubated in RPMI-1640 supplemented with 10% FBS and 2 mM L-glutamine with PHA-L (1-2 $\mu\text{g/ml}$) (Sigma), 10×10^6 irradiated feeder cells (CESS), rhIL-2 (200 IU/ml) (Proleukin, Chiron, Emeryville, CA), and IL-4 (10 ng/ml) (R&D Systems, Minneapolis, MN). T cell clones were usually allowed to expand for 10-14 days before re-stimulation. A panel of antigen non-specific T cell clones was examined for the expression of CD70. Table 2 summarizes the characteristics of the T cell clones examined. This panel contains both CD4^+ and CD8^+ T cells clones that appear to belong to different T cell differentiation pathways as suggested by their cytokine profiles. Significant levels of CD70 were detectable on all of the clones examined. T cell clones were activated by PHA, irradiated feeder cells (CESS), and rIL-2 and the expression of CD25 and CD70 was monitored by flow cytometric analysis. Clone C1A, representative of other T cell clones, showed extensive upregulation of CD25, which peaked on day 2 (Figure 13). This is indicative of T lymphocyte activation. Expression of CD25 gradually declined in the following days. The induction of CD70 expression paralleled that of CD25. The peak CD70 induction was also observed after 2 days of stimulation. Considerable CD70 expression was still detectable on day 8. All T cell clones examined, including those listed in Table 2, showed similar kinetics and magnitudes of activation-induced expression of CD70.

Table 2: T cell clone phenotypes and activation-induced expression of CD70

T cell clone	Phenotype			
	T-lineage marker	CD70 expression*	Cytokine profile	Designation
3.27.2	CD3, CD4	15	IL-4, IL5, IL-13, IFN γ	Th0
4.01.1	CD3, CD4	7	IL-4, IL5, IL-13, IFN γ	Th0
20G5	CD3, CD4	3	IL-4, IL-13	Th2
40D8	CD3, CD4	8	IL-4, IL-13	Th2
C1A	CD3, CD8	3	IL-4, IL-13, IFN γ	Tc0
C2A	CD3, CD8	12	IL-4, IL-13, IFN γ	Tc0

* Ratio between the mean log fluorescence intensities for anti-CD70 binding and control IgG binding

5

Example 10: Proliferation Inhibitory Effects of Anti-CD70 ADCs on Activated T Cells

[0314] The effects of 1F6 ADCs on the proliferation of activated T cells were examined using activated T cell clones, mixed lymphocyte reactions (MLR), and antigen-primed T cells.

- 10 [0315] Standard proliferation assays in 96-well format in quadruplicates were used to evaluate the effects of anti-CD70 ADCs on activated T cells. For MLR, PBMCs were seeded at 50,000 cells/well with 50,000 irradiated allogeneic CESS cells in a total volume of 200 μ l of medium. For tetanus toxoid-induced T cell proliferation, tetanus toxoid-reactive T cells were seeded at 20,000 cells/well with 2,000 autologous DCs in a total volume of 200 μ l.
- 15 Dialyzed and clarified tetanus toxoid (Colorado Serum Company, Denver, CO) was used at a final concentration of 1:50 dilution. Two-day activated T cell clones were seeded at 10,000 cells per well. Proliferation assays was usually carried out for 72 to 120 hours. Tritiated thymidine (3 H-TdR) incorporation during the last 16 hours of incubation and scintillation counting were used to assess DNA synthesis. For analysis, the amount of thymidine
- 20 incorporated by the treated cells was compared to that of the untreated control cells.

- [0316] In the first system, the responses of T cell clones to 1F6 ADCs were examined. Resting T cell clones were activated to induce CD70 expression as described in Figure 13. After 2 days of activation, a portion of the cells was analyzed for CD25 and CD70 expression by flow cytometry to confirm cellular activation and CD70 induction. The remaining cells
- 25 were pelleted and resuspended in new medium containing 200 IU/ml of rIL-2 or 200 IU/ml of IL-2 and 10 ng/ml IL-4. Cells were then plated out at 10,000 cells/well in a final volume of 200 μ l of medium containing graded concentrations of 1F6 ADCs or the non-binding IgG ADCs. Cells were incubated for an additional 72 hours with the last 16 hours pulsed with 3 H-TdR to assess cellular DNA synthesis. Results for the responses of three T cell clones toward
- 30 ADC treatment was shown in Figure 14. 1F6 ADCs at concentrations higher than 0.01 μ g/ml significantly inhibited the proliferation of the T cell clones. 1F6-vcAFP appeared to be more active than 1F6-vcMMAE; the IC₅₀s for 1F6-vcAFP on all 3 clones were found to be below 0.05 μ g/ml. The control ADC cIgG-vcMMAE and cIgG-vcAFP did not significantly inhibit proliferation at concentrations below 2 μ g/ml, confirming the antigen specificity of the 1F6
- 35 ADCs.

5 [0317] In the second system, an MLR was set up between PBMC and an irradiated allogeneic stimulator CESS. Allogeneic activation of peripheral blood T cells was achieved by conventional mixed lymphocyte reactions (MLR). MLR were initiated by mixing PBMCs and irradiated allogeneic Epstein-Barr virus (EBV)-transformed B lymphoblastoid CESS cells (ATCC, Manassas, VA) at a ratio of one T cell to ten irradiated CESS cells. T cell
10 density was adjusted to 0.25×10^6 T cells/ml of RPMI-1640 supplemented with 10% FBS and 2 mM L-glutamine. Rapid proliferation and expansion of alloreactive T cells was observed after approximately 96 hours of culture. After 5 days of stimulation, cells were assessed for CD70 expression using the anti-CD70 mAb 1F6 by flow cytometry. Flow cytometric analysis showed that most of the cells within the viable gate were CD3⁺ T cells and that
15 CD70 was expressed on the viable CD3⁺ T cells. Figure 15, upper panel, shows the expression of CD70 at the end of the MLR (120 hours after initiation). In parallel cultures in 96-well plates, graded doses of either 1F6-vcMMAE or 1F6-vcAFP were included in the cultures at initiation. DNA synthesis was assayed during the last 16 hours of the 5-day culture by a pulse of ³H-TdR. Both 1F6-vcMMAE and 1F6-vcAFP substantially inhibited T
20 cell proliferation (Figure 15, lower panel). 1F6-vcAFP was more active than 1F6-vcMMAE, with an IC₅₀ of approximately 0.1 µg/ml.

[0318] In the third system, the effect of 1F6 and 1F6 ADCs on antigen-induced T cell proliferation was examined. To evaluate antigen-specific T cell activation, T cells specific against tetanus toxoid were enriched from PBMC by multiple rounds of antigenic stimulation
25 with tetanus toxoid in the presence of autologous DCs. Before antigenic stimulation of T cells, mature autologous DCs pulsed with tetanus toxoid were prepared as described above. Autologous PBMC were then co-cultured with antigen-pulsed mature DCs at a ratio of 10 PBMC to 1 DC. At the density of $0.25\text{--}0.5 \times 10^6$ PBMCs/ml of RPMI-1640 supplemented with 10% FBS and 2 mM L-glutamine. T cell activation and expansion were allowed to
30 continue for 7 days. Viable T cells were harvested from the culture and re-stimulated with tetanus toxoid-pulsed autologous mature DCs again under conditions identical to those in first round of activation. Two additional, identical rounds of activation were conducted to further enrich for tetanus toxoid-specific T cells.

[0319] Tetanus toxoid-specific T cells were enriched and expanded by four consecutive
35 rounds of stimulation by tetanus toxoid and autologous DCs as described above. After the 4th round of stimulation, resting tetanus toxoid-specific T cells were plated in 96-well plates at 20,000 cells/well in the presence of autologous DCs (at 2,000 cells/well) and tetanus

- 5 toxoid. Graded doses of 1F6, 1F6-vcMMAE, and 1F6-vcAFP were also included in the cultures at the initiation. DNA synthesis was assayed during the last 16 hours by ^3H -TdR incorporation. Unconjugated 1F6 did not show a significant effect on tetanus toxoid-induced T cell proliferation, whereas both 1F6-vcMMAE and 1F6-vcAFP exerted substantial inhibition at doses higher than 0.03 $\mu\text{g/ml}$ (Figure 16).

10

Example 11: CD70 Expression on Activated T cells during an Antigen-Specific *In Vitro* Immune Response

- [0320] A 9-amino acid peptide (GILGFVFTL, M1 peptide) derived from the influenza virus matrix protein binds to the peptide-binding groove of the HLA-A0201 molecule.
- 15 Presentation of the M1 peptide by HLA-A0201 expressing antigen presenting cells to autologous T cells specifically stimulates the activation and expansion of CD8^+ cytotoxic T cells expressing the T cell receptor $\text{V}\beta 17$ chain (Lehner *et al.*, 1995, *J. Exp. Med.* 181: 79-91), constituting a convenient *in vitro* experimental system to track the activation and expansion of antigen-specific T cells to their cognate antigen.
- 20 [0321] To examine CD70 expression on activated antigen-specific T cells, PBMCs from a normal donor expressing HLA-A0201 were stimulated with the M1 peptide. PBMCs were seeded at 2×10^6 cells/ml with 5 $\mu\text{g/ml}$ of M1 peptide in AIMV medium supplemented with 5% human AB serum. IL-2 (Proleukin, Chiron) and IL-15 (R&D Systems, MN) were added to final concentration of 20 IU/ml and 5 ng/ml, respectively, once every two days beginning
- 25 on day 2 after culture initiation. The expansion of $\text{CD8}^+/\text{V}\beta 17^+$ T cells and induction of CD70 on the $\text{CD8}^+/\text{V}\beta 17^+$ was followed by three-color flow cytometry. $\text{V}\beta 17^+$ T cells were identified by the anti-TCRV $\beta 17$ mAb clone E17.5F3 (Beckman Coulter, Miami, FL). Results from a representative experiment are shown in Figures 17A and B. Figure 17A, upper panels, shows that only 0.9% of the cells within the lymphocyte population were
- 30 $\text{CD8}^+/\text{V}\beta 17^+$ two days after culture initiation. T cell expansion was only evident within the $\text{CD8}^+/\text{V}\beta 17^+$ population. The percentage of $\text{CD8}^+/\text{V}\beta 17^+$ progressively increased to 23% on day 11. CD70 expression became detectable 3 days after antigen stimulation (Figure 17A, lower panel). On day 7, approximately 60% of the expanding $\text{CD8}^+/\text{V}\beta 17^+$ cells expressed CD70 (Figure 17A, lower left panel). The highest level of CD70 expression as indicated by
- 35 the mean fluorescence intensity (MFI) was also detected on day 7 (Figure 17A, lower right

5 panel). The percentage of CD70⁺/CD8⁺/Vβ17⁺ cells and the MFI of CD70 expression on the CD8⁺/Vβ17⁺ started to decline thereafter. Whereas CD70 was clearly expressed on the CD8⁺/Vβ17⁺ cells, no CD70 could be detected on the CD8⁺/Vβ17⁻ cells (Figure 17B). These results confirmed that CD70 induction was restricted to the activated T cells responding to the antigenic stimulation but not the bystander, antigen non-specific T cells.

10

Example 12: *In Vitro* Deletion of CD70⁺ Antigen-Specific T Cells by Anti-CD70 ADCs

[0322] PBMCs from a normal donor expressing HLA-A0201 were stimulated with the M1 peptide as described in Example 11. On day 5, cells were harvested, washed, and re-seeded at 0.5-1 x 10⁶ cells/ml (approximately 100,000 of CD8⁺/Vβ17⁺ T cells) in fresh AIMV
15 medium supplemented with 5% human AB, 20 IU/ml of IL-2, and 5 ng/ml of IL-15. 1F6 or control non-binding IgG (cIgG) ADCs were added to some cultures to a final concentration of 1 μg/ml. Total viable cell counts were conducted 24, 48, and 76 hours after ADC addition. Two-color flow cytometry was conducted to determine the percentages of CD8⁺/Vβ17⁺ cells among the viable cells. The absolute number of CD8⁺/Vβ17⁺ T cells in each culture was then
20 calculated. The number of CD8⁺/Vβ17⁺ T cells in the two control cultures increased to more than 800,000 after 76 hours (Figure 18, upper panel). Substantial inhibition of CD8⁺/Vβ17⁺ T cell expansion was seen in the 1F6 ADC-treated cultures. The number of CD8⁺/Vβ17⁺ T cells was approximately 400,000 in the 1F6-vcMMAE-treated culture, compared to >800,000 in the cIgG-vcMMAE-treated culture. Virtually no expansion of CD8⁺/Vβ17⁺ T cells
25 occurred in the 1F6-vcAFP-treated culture, whereas >500,000 of CD8⁺/Vβ17⁺ T cells were present in the cIgG-vcAFP-treated culture. The effects of 1F6 ADCs on the CD8⁺/Vβ17⁻ cells were also evaluated. Figure 18, lower panel, shows that the CD8⁺/Vβ17⁻ cells in the 1F6-vcAFP- and 1F6-vcMMAE-treated cultures were <40% and <20%, respectively, of the control untreated culture at the end of the experiment. In contrast, the antigen non-specific
30 bystander, antigen non-specific CD8⁺/Vβ17⁻ cells in the ADC-treated cultures were >60% of the untreated culture. These results demonstrate that 1F6 ADCs specifically targeted the CD70⁺/CD8⁺/Vβ17⁺ T cells and exerted limited effects on the CD70⁻/CD8⁺/Vβ17⁺ T cells that co-inhabited the same culture, suggesting limited bystander toxicity of the 1F6 ADCs.

[0323] The potency of three 1F6 ADCs was then compared. HLA-A0201 PBMCs were
35 stimulated with the M1 peptide and re-seeded as described for Figure 18 with the various

5 ADCs at the indicated concentrations (Figure 19). The percent of CD8⁺/Vβ17⁺ T cells after a 96-hour incubation was determined. A dose-dependent inhibition of CD8⁺/Vβ17⁺ T cell expansion was observed with 1F6-vcMMAE, -vcAFP, and -vcMMAF, while the corresponding IgG control ADCs at the top concentration of 1 μg/ml showed no detectable inhibitory activity compared to the no drug control culture. The IC₅₀ values for 1F6-vcAFP
10 and 1F6-vcMMAF were between 0.1 and 0.01 μg/ml. The inhibitory activity of 1F6-vcMMAE appeared to be weaker than both 1F6-vcAFP and 1F6-vcMMAF.

[0324] A re-stimulation assay as used to evaluate the functional capacity of the bystander, antigen non-specific CD8⁺/Vβ17⁺ T cells. HLA-A0201 expressing PBMCs were stimulated with the M1 peptide and then treated with 1 μg/ml of 1F6-vcMMAF on day 5 to delete the
15 CD70⁺/CD8⁺/Vβ17⁺ T cells as described in Figures 18 and 19. As expected the number of CD8⁺/Vβ17⁺ T cells in the ADC-treated culture was lower than the untreated culture, 14% versus 47% (Figure 20, left column). Responses of the remaining cells to a mitogenic re-stimulation were assessed. In order to normalize for the number of Vβ17⁺ T cells used in the re-stimulation assay and minimize the contribution of CD8⁺/Vβ17⁺ T cells to the proliferation
20 response, Vβ17⁺ T cells were enriched by negative immunoselection. Briefly, day 9 cells were harvested, washed, and resuspended to 5 x 10⁷ cells/ml in medium containing an anti-TCRVβ17 mAb (clone E17.5F3) at 1-2 μg/10⁶ target T cells. After incubation at 4°C for 30 minutes, unbound anti-TCRVβ17 mAb was removed by washing the cells twice with PBS containing 0.1% human serum. The cells were resuspended at a concentration of 2-5 x
25 10⁶/ml two washes in medium and Dynabeads M-450 goat anti-mouse IgG paramagnetic beads (DynaL Biotech Inc., Lake Success, NY) were added at a ratio of 4 paramagnetic beads to 1 target T cell. The mixture was rotated at 4°C for 30 minutes. Paramagnetic beads with bound Vβ17⁺ T cells were then separated from the rest of the cells by a magnetic device. Removal of Vβ17⁺ T cells was confirmed by two-color flow cytometry. Figure 20, middle
30 column, shows that only 3% and 5% of the cells were left in the 1F6-vcMMAF-treated and untreated cultures, respectively. Anti-CD3 and anti-CD28 immobilized onto tissue culture wells at graded concentrations were used to re-stimulate the Vβ17⁺ cells in the absence of any exogenous IL-2. Twenty thousand Vβ17⁺ cells were seeded per well in 96-well plates in a total of 200 μl of AIMV medium supplemented with 5% human serum. Cells were cultured
35 for of 96 hours, and DNA synthesis was assayed by ³H-TdR incorporation during the last 18 hours of culture. Vβ17⁺ T cells enriched from the 1F6-vcMMAF-treated culture

5 demonstrated a dose-dependent response to anti-CD3 plus anti-CD28 re-stimulation comparable to that of the untreated cells. This suggests that, although the V β 17⁺ T cells were incubated with the CD8⁺/V β 17⁺ cells in the presence of 1F6-vcMMAF for 4 days, they had retained their proliferation capacity. More importantly, their ability to mount a proliferation response in the absence of any exogenous IL-2 illustrates that they were also functionally
 10 intact with regard to their ability to express and secrete the necessary cytokine(s) needed for their proliferation. Taken together, these results suggest that anti-CD70 ADCs can selectively delete CD70⁺ activated T cells without causing substantial collateral damage to bystander T cells. This characteristic suggests that they can be applied as immunosuppressive agents that may have minimum impact on the immune repertoire of the
 15 host.

Example 13: Treatment of Experimental Allergic Encephalomyelitis by Administration of Anti-CD70 ADCs

[0325] Studies indicate a role for CD70/CD27-mediated T cell-T cell interactions in enhancing the Th₁-mediated immune responses in cell-mediated autoimmune diseases, including, for example, autoimmune demyelinating diseases. In this example, experimental
 20 allergic encephalomyelitis (EAE), an animal model of the demylelinating disease multiple sclerosis (MS), is treated with an ADC comprising an antibody that (a) is conjugated to AFP or MMAE with a val-cit (vc) linker and (b) recognizes an epitope of murine CD70
 25 corresponding the 1F6 epitope of human CD70.

[0326] *Induction and clinical assessment of Experimental allergic encephalomyelitis (EAE):* R-EAE (relapsing EAE) is induced in six- to seven-week-old female SJL mice by subcutaneous immunization with 100 μ l of complete Freund's adjuvant (CFA) emulsion containing 200 μ g of *Mycobacterium tuberculosis* H37Ra and 40 μ g of the immunodominant
 30 epitope of proteolipid protein, PLP₁₃₉₋₁₅₁. The signs of EAE are scored on a 0 to 5 scale as follows: (0) normal; (1) limp tail or hind limb weakness; (2) limp tail and hind-limb weakness (waddling gait); (3) partial hind-limb paralysis; (4) complete hind-limb paralysis; and (5) moribund. A relapse is defined as a sustained increase (more than 2 days) in at least one full grade in clinical score after the animal had improved previously at least a full clinical
 35 score and had stabilized for at least 2 days. The data are plotted as the mean clinical score for

5 all animals in a particular treatment group or as the relapse rate (total number of relapses in a group divided by the total number of mice in that group).

[0327] *Anti-CD70 ADC Administration Regimens:* Anti-CD70 ADC (0.1-3 mg/kg body weight) is administered intraperitoneally in a total volume of 100 μ l. Mice are treated 3 times per week for 3 consecutive weeks (9 total treatments). Treatment is initiated before disease onset (day 7) or at the peak of acute disease (day 14). As a control, one group of EAE-
10 induced mice are left untreated.

[0328] *Inhibition of TNF- α and IFN- γ induction:* The demonstration of induction of TNF- α and IFN- γ in brains of EAE shows an inflammatory disease process indicative of EAE disease progression and inhibition of these cytokines in brains of SJL mice treated with anti-
15 CD70 ADC indicates the value of anti-CD70 ADC therapy in preventing or treating EAE. Brains are obtained from from at least three animals treated preclinically (at day 13, after three treatments, and day 26, after nine treatments) and at peak of acute disease (at day 20, after three treatments, and day 33, after nine treatments). Brains are fixed (10% buffered formalin), and tissues are embedded in paraffin and sectioned. Sections are then
20 independently stained for TNF- α or IFN- γ by incubation with a primary antibody specific for the respective cytokine, followed by incubation with a secondary antibody conjugated to FITC. Tissue sections are then mounted in mounting media and analyzed by immunofluorescence microscopy. Decreased levels of TNF- α or INF- γ staining in ADC-treated mice versus the untreated EAE-induced mice shows inhibition of inflammatory
25 cytokine induction using anti-CD70 ADC therapy.

[0329] *Inhibition of Disease Symptoms or Relapse Rates:* EAE-induced SJL mice in the treatment group are compared with untreated EAE-induced mice to assess the efficacy of anti-CD70 ADC therapy in either preventing disease onset or treating established disease. For mice treated preclinically, a decrease in the mean score for EAE disease, as compared to
30 the untreated control group, demonstrates the efficacy of anti-CD70 ADC therapy in preventing disease. For mice treated at the peak of acute disease, either (a) a decrease in the relapse rate or (b) a decrease in the post-treatment mean score for EAE, as compared to the untreated control group, demonstrates the efficacy of anti-CD70 ADC therapy in treating established disease.

35

- 5 **[0330]** The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.
- 10 **[0331]** Various references, including patent applications, patents, and scientific publications, are cited herein, the disclosures of each of which is incorporated herein by reference in its entirety.

WHAT IS CLAIMED IS:

1. A method for the treatment of a CD70-expressing cancer in a subject, the method comprising:
administering to the subject, in an amount effective for the treatment, an antibody-drug conjugate comprising an antibody that binds to CD70 and wherein the drug is a cytotoxic or cytostatic agent.
2. The method of claim 1, wherein the antibody competes for binding to CD70 with monoclonal antibody 1F6 or 2F2.
3. The method of claim 2, wherein the antibody comprises at least one polypeptide region selected from the group consisting of
 - (a) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:6;
 - (b) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:8;
 - (c) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:10;
 - (d) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:16;
 - (e) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:18;
 - (f) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:20;
 - (g) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:26;
 - (h) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:28;
 - (i) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:30;
 - (j) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:36;
 - (k) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:38; and

- (l) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:40.
4. The method of claim 3, wherein the antibody comprises the H1, H2, and H3 regions of (a), (b), and (c); or the H1, H2, and H3 regions of (g), (h), and (i).
5. The method of claim 4, wherein the polypeptide regions of (a), (b), (c), (d), (e), and (f) have, respectively, the amino acid sequences set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10; SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20.
6. The method of claim 4, wherein the antibody comprising the H1, H2, and H3 regions of (a), (b), and (c) further comprises the L1, L2, and L3 regions of (d), (e), and (f); or the antibody comprising the H1, H2, and H3 regions of (g), (h), and (i) further comprises the L1, L2, and L3 regions of (j), (k), and (l).
7. The method of claim 4, wherein the antibody comprises a heavy chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO 22.
8. The method of claim 7, wherein the heavy chain variable region has the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:22.
9. The method of claim 7, wherein the antibody, comprising the heavy chain variable region having at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:22, further comprises, respectively, a light chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.
10. The method of claim 9, wherein the light chain variable region has the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.
11. The method of claim 10, wherein the antibody is monoclonal antibody 1F6 or 2F2.

12. The method of claim 11, wherein the antibody-drug conjugate is selected from the group consisting of 1F6-val-cit-AFP, 1F6-val-cit-MMAF, 1F6-val-cit-MMAE, 2F2-val-cit-AFP, 2F2-val-cit-MMAF, and 2F2-val-cit-MMAE.
13. The method of claim 1, wherein the antibody is a chimeric, humanized, or human antibody.
14. The method of claim 13, wherein the chimeric antibody comprises a human constant region.
15. The method of claim 1, wherein the antibody is multivalent.
16. The method of claim 1, wherein the cytotoxic agent is selected from the group consisting of an auristatin, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a duocarmycin, a taxane, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid.
17. The method of claim 1, wherein the cytotoxic agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, paclitaxel, docetaxel, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, chaliceamicin, maytansine, DM-1, or netropsin.
18. The method of claim 1, wherein the cytotoxic agent is an anti-tubulin agent.
19. The method of claim 18, wherein the anti-tubulin agent is an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin.
20. The method of claim 18, wherein the antitubulin agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcimid, estramustine, cemadotin, discodermolide, maytansine, DM-1, or eleutherobin.
21. The method of claim 20, wherein the cytotoxic agent is AFP, MMAF, or MMAE.

22. The method of claim 1, wherein the antibody is conjugated to the cytotoxic agent via a linker.
23. The method of claim 22, wherein the linker is cleavable under intracellular conditions.
24. The method of claim 23, wherein the cleavable linker is a peptide linker cleavable by an intracellular protease.
25. The method of claim 24, wherein the peptide linker is a dipeptide linker.
26. The method of claim 25, wherein the dipeptide linker is a val-cit linker or a phe-lys linker.
27. The method of claim 23, wherein the cleavable linker is hydrolyzable at a pH of less than 5.5.
28. The method of claim 27, wherein the hydrolyzable linker is a hydrazone linker.
29. The method of claim 23, wherein the cleavable linker is a disulfide linker.
30. The method of claim 1, wherein the CD70-expressing cancer is selected from the group consisting of a kidney tumor, a B cell lymphoma, a thymic carcinoma, a nasopharyngeal carcinoma, and a brain tumor.
31. The method of claim 30, wherein the kidney tumor is a renal cell carcinoma.
32. The method of claim 30, wherein the brain tumor is a glioma, a glioblastoma, or a meningioma.
33. The method of claim 1, wherein the subject is human.
34. A method for the treatment of an immunological disorder in a subject, the method comprising:

administering to the subject, in an amount effective for the treatment, an antibody-drug conjugate comprising an antibody that binds CD70 and wherein the antibody is conjugated to a cytotoxic agent or an immunosuppressive agent.

35. The method of claim 34, wherein the antibody competes for binding to CD70 with monoclonal antibody 1F6 or 2F2.

36. The method of claim 35, wherein the antibody comprises at least one polypeptide region selected from the group consisting of

- (a) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:6;
- (b) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:8;
- (c) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:10;
- (d) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:16;
- (e) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:18;
- (f) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:20;
- (g) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:26;
- (h) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:28;
- (i) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:30;
- (j) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:36;
- (k) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:38; and
- (l) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:40.

37. The method of claim 36, wherein the antibody comprises

the H1, H2, and H3 regions of (a), (b), and (c); or
the H1, H2, and H3 regions of (g), (h), and (i).

38. The method of claim 37, wherein the polypeptide regions of (a), (b), (c), (d), (e), and (f) have, respectively, the amino acid sequences set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10; SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20.

39. The method of claim 37, wherein
the antibody comprising the H1, H2, and H3 regions of (a), (b), and (c) further comprises the L1, L2, and L3 regions of (d), (e), and (f); or
the antibody comprising the H1, H2, and H3 regions of (g), (h), and (i) further comprises the L1, L2, and L3 regions of (j), (k), and (l).

40. The method of claim 37, wherein the antibody comprises a heavy chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO 22.

41. The method of claim 40, wherein the heavy chain variable region has the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:22.

42. The method of claim 40, wherein the antibody, comprising the heavy chain variable region having at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:22, further comprises, respectively, a light chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

43. The method of claim 42, wherein the light chain variable region has the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

44. The method of claim 43, wherein the antibody is monoclonal antibody 1F6 or 2F2.

45. The method of claim 44, wherein the antibody-drug conjugate is selected from the group consisting of 1F6-val-cit-AFP, 1F6-val-cit-MMAF, 1F6-val-cit-MMAE, 2F2-val-cit-AFP, 2F2-val-cit-MMAF, and 2F2-val-cit-MMAE.

46. The method of claim 34, wherein the antibody is a chimeric, humanized, or human antibody.

47. The method of claim 46, wherein the chimeric antibody comprises a human constant region.

48. The method of claim 34, wherein the antibody is multivalent.

49. The method of claim 34, wherein the cytotoxic agent is selected from the group consisting of an auristatin, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a duocarmycin, a taxane, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid.

50. The method of claim 34, wherein the cytotoxic agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, paclitaxel, docetaxel, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, chaliceamicin, maytansine, DM-1, or netropsin.

51. The method of claim 34, wherein the cytotoxic agent is an anti-tubulin agent.

52. The method of claim 51, wherein the anti-tubulin agent is an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin.

53. The method of claim 52, wherein the antitubulin agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcemid, estramustine, cemadotin, discodermolide, maytansine, DM-1, or eleutherobin.

54. The method of claim 53, wherein the cytotoxic agent is AFP, MMAF, or MMAE.

55. The method of claim 34, wherein the immunosuppressive agent is gancyclovir, etanercept, cyclosporine, tacrolimus, rapamycin, cyclophosphamide, azathioprine, mycophenolate mofetil, methotrexate, cortisol, aldosterone, dexamethasone, a cyclooxygenase inhibitor, a 5-lipoxygenase inhibitor, or a leukotriene receptor antagonist.

56. The method of claim 34, wherein the antibody is conjugated to the cytotoxic or immunosuppressive agent via a linker.

57. The method of claim 56, wherein the linker is cleavable under intracellular conditions.
58. The method of claim 57, wherein the cleavable linker is a peptide linker cleavable by an intracellular protease.
59. The method of claim 58, wherein the intracellular protease is a lysosomal protease or an endosomal protease.
60. The method of claim 58, wherein the peptide linker is a dipeptide linker.
61. The method of claim 60, wherein the dipeptide linker is a val-cit linker or a phe-lys linker.
62. The method of claim 57, wherein the cleavable linker is hydrolyzable at a pH of less than 5.5.
63. The method of claim 62, wherein the hydrolyzable linker is a hydrazone linker.
64. The method of claim 57, wherein the cleavable linker is a disulfide linker.
65. The method of claim 34, wherein the immunological disorder is a T cell-mediated immunological disorder.
66. The method of claim 65, wherein the T cell mediated immunological disorder is a T-cell mediated and include activated T cell expressing CD70.
67. The method of claim 66, wherein resting T cells are not substantially depleted by administration of the antibody-drug conjugate.
68. The method of claim 65, wherein the T cell-mediated immunological disorder is rheumatoid arthritis, multiple sclerosis, psoriasis, Sjorgren's syndrome, Hashimoto's thyroiditis, Grave's disease, primary biliary cirrhosis, Wegener's granulomatosis, tuberculosis, or acute graft versus host disease.

69. The method of claim 34, wherein the immunological disorder is an activated B-lymphocyte disorder.

70. The method of claim 34, wherein the subject is human.

71. An antibody-drug conjugate comprising:
an antibody that binds to CD70 and is conjugated to a cytotoxic agent or an immunosuppressive agent, wherein the antibody-drug conjugate exerts (a) a cytotoxic or cytostatic effect on a CD70-expressing cancer cell line or (b) a cytotoxic, cytostatic, or immunosuppressive effect on a CD70-expressing immune cell.

72. The antibody-drug conjugate of claim 71, wherein the antibody competes for binding to CD70 with monoclonal antibody 1F6 or 2F2.

73. The antibody-drug conjugate of claim 72, wherein the antibody comprises at least one polypeptide region selected from the group consisting of

- (a) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:6;
- (b) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:8;
- (c) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:10;
- (d) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:16;
- (e) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:18;
- (f) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:20;
- (g) an H1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:26;
- (h) an H2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:28;
- (i) an H3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:30;

- (j) an L1 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:36;
- (k) an L2 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:38; and
- (l) an L3 region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:40.

74. The antibody-drug conjugate of claim 73, wherein the antibody comprises

the H1, H2, and H3 regions of (a), (b), and (c); or
the H1, H2, and H3 regions of (g), (h), and (i).

75. The antibody-drug conjugate of claim 74, wherein the polypeptide regions of (a), (b), (c), (d), (e), and (f) have, respectively, the amino acid sequences set forth in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10; SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20.

76. The antibody-drug conjugate of claim 74, wherein
the antibody comprising the H1, H2, and H3 regions of (a), (b), and (c) further comprises the L1, L2, and L3 regions of (d), (e), and (f); or
the antibody comprising the H1, H2, and H3 regions of (g), (h), and (i) further comprises the L1, L2, and L3 regions of (j), (k), and (l).

77. The antibody-drug conjugate of claim 74, wherein the antibody comprises a heavy chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO 22.

78. The antibody-drug conjugate of claim 77, wherein the heavy chain variable region has the amino acid sequence set forth in SEQ ID NO:2 or SEQ ID NO:22.

79. The antibody-drug conjugate of claim 77, wherein the antibody, comprising the heavy chain variable region having at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:22, further comprises, respectively, a light chain variable region having at least 80% sequence identity to the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

80. The antibody-drug conjugate of claim 79, wherein the light chain variable region has the amino acid sequence set forth in SEQ ID NO:12 or SEQ ID NO:32.

81. The antibody-drug conjugate of claim 80, wherein the antibody is monoclonal antibody 1F6 or 2F2.

82. The antibody-drug conjugate of claim 81, wherein the antibody-drug conjugate is selected from the group consisting of 1F6-val-cit-AFP, 1F6-val-cit-MMAF, 1F6-val-cit-MMAE, 2F2-val-cit-AFP, 2F2-val-cit-MMAF, and 2F2-val-cit-MMAE.

83. The antibody-drug conjugate of claim 71, wherein the antibody is a human, humanized, or chimeric antibody.

84. The antibody-drug conjugate of claim 71, wherein the antibody is multivalent.

85. The antibody-drug conjugate of claim 71, wherein the cytotoxic agent is selected from the group consisting of an auristatin, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a duocarmycin, a taxane, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid.

86. The antibody-drug conjugate of claim 71, wherein the cytotoxic agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, paclitaxel, docetaxel, CC-1065, SN-38, topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, chaliceamicin, maytansine, DM-1, or netropsin.

87. The antibody-drug conjugate of claim 71, wherein the cytotoxic agent is an anti-tubulin agent.

88. The antibody-drug conjugate of claim 87, wherein the anti-tubulin agent is an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin.

89. The antibody-drug conjugate of claim 86, wherein the antitubulin agent is AFP, MMAF, MMAE, AEB, AEVB, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B,

WO 2004/073656

PCT/US2004/005247

WO 2004/073656

"

PCT/US2004/005247

FIGURE 1**IE6 VL nucleotide sequence:**

```

1  atggagacag acacactct gttatgggta ctctgctct gggttcagg ttccactggt 60
61  gacattgtgc tgacacagtc tctgtctcc ttactgttat ctctgggca gagggccacc 120
121  atctcatgca gggccagcaa aagtgtcagt acatctggct atagtatat gcatgggtat 180
181  caacagaaac caggacagcc acccaactc ctcatctat ttgcatpcaa cctagaatct 240
241  ggggtccctg ccaggttcag tggcagtggt ttgggacag acttcacct caacatccat 300
301  cctgtggagg aggaggatgc tgcacacct tatctgtcagc acagtaggga gggtccgtgg 360
361  acgttcggtg gaggcaccaa gctggaaatc aaacgg 396

```

IE6 VL amino acid sequence:

```

-20 MESTYFLLMV LLINVPQSTG
1  DIVLTQSPAS LANSIGURAT ISCRASKSVS YSGISFMHW QKRPGRPKL IYLAQMLES 60
61  GVPARFSGSG SGTDFTLNHH FVEEDAAFY YQHSREVPW TEGGTKLEI KR 112

```

IE6 VH nucleotide sequence:

```

1  atggccttggg ttgtggacctt yctattcttg atggcagctg cccaaagtgc ccaagcacag 60
61  atccagttgg ttgagttctgg acctgaggtg aagaagcctg gagagacagt caagatctcc 120
121  tgcagggttt ctgggtatc ctctcaaac tatggaatga actgggtgaa gcaggtcca 180
181  ggaagggttt taagttgat gggctggata aacacclaca ctggagagcc aacataigct 240
241  gatgccctta agggacaggt ttgccttctt ttggaacct ctgccagcac tgcctatttg 300
301  cagatcaaca acctcaaaa tggggacacg gctacatat tctgtgcaag agactacggc 360
361  gactatggtg ttgactactg ggttcaagga acctcagtca cagtcctctc a 411

```

IE6 VH amino acid sequence:

```

-19 MRAVATLLFL IAAQASAG
1  QIQLVDSGSE VKKPGTIVIA SCASGIYFT KGENAVROA PGRGLKRW EMTTCEPTY 60
61  ADAFKERFAF SLETSASTAY LQINLKQED TATFTCARDY GDYGADYNGQ GTSVTVSS 118

```

2/23

FIGURE 2

2F2 VL nucleotide sequence:

```

1  atgagacag acacactcct gttatggta ctgctgctct gggttccagg ttccactggt 60
61  gacatgtgc tgacacagtc tctgtcttc ttaactgtat ctctggggca gaagaccacc 120
121 atctcatgca gggccagcaa gagtgtcagt acatctggct atagtcttat gcactggtae 180
181 caactgaac caggacagtc acccaaacct ctcactctat ttggtccaa cttaccatct 240
241 ggggtccctg ccaggttcag aggcagtggg tctggggcag acttaccct caaatccat 300
301 cctgtggagg aggcagatgc tgcaacctat tactgtcagc acagtaggga gttccgtac 360
361 acgttcggag gggggaccaa gctggaaata acacgg 396

```

2F2 VL amino acid sequence:

```

0
-20 METDTLEIMV LLLNVPGSTG
1  DIVLTQSPAS LTVALGQRTT ISCRASKSVS TSGYSEHWI QLRPGSPKL LYLASNLPS 60
61  GVPAKPSGSG SGTDFTLKH FVEEDDAATY YCHSRREPY TFSGGTALET YR 112

```

2F2 VH nucleotide sequence:

```

1  atggaatgga cctgggtctt tctcttctc ctgtcagtaa ctgcagatgt ccaatccag 60
61  gttcagctgc aacagtctgg aactgagctg atgacgcccg gggctcagt gacgatgtcc 120
121 tgcaagactt ctggctacac attcagtagc tactggatag agtgggtaaa acagaggcct 180
181 ggacatggcc ttgagtggat tggagaatct ttaccctgaa gtgttatfac tgactacaat 240
241 gagaagtcca aggccaaagg cacattcact gcagatacat cctcaaacac agcttaccatg 300
301 caactcagca ggcctggcatc tggaggactct gcgtctatt actgtgcaag atgggtagg 360
361 ctctatgcta tggactactg gggtaacagg acctcagtca ccgtatccctc a 411

```

2F2 VH amino acid sequence:

```

0
-19 METWVLEL LSVTADVQS
1  QVQLQSGTE LTTCASVTH SCKTSGITFS TWIENAKQK PCHLEWIGK ILGPSGITDY 60
61  NERFKAKATF TADYSSNTAY LQLSLASED SNTYCAKND RLRYMDYWGQ GTSVTVSS 118

```

3/23

FIGURE 3

1F6 CDR-I1 (residues 23 to 38): R A S K S V S T S G Y S F M H
 2F2 CDR-I1 (residues 23 to 38): R A S K S V S T S G Y S F M H

1F6 CDR-I2 (residues 54 to 60): L A S N L E S
 2F2 CDR-I2 (residues 54 to 60): L A S N L P S

1F6 CDR-I3 (residues 93 to 101): Q H S R E V P N T
 2F2 CDR-I3 (residues 93 to 101): Q H S R E I P Y T

1F6 CDR-H1 (residues 26 to 35): G Y T F T N Y G M N
 2F2 CDR-H1 (residues 26 to 35): G Y T F S T Y N I E

1F6 CDR-H2 (residues 49 to 56): N I N T Y T G E P T Y A D A F K G
 2F2 CDR-H2 (residues 49 to 56): E I L G P S G Y T D Y N E K F K A

1F6 CDR-H3 (residues 99 to 107): D Y G D Y G M D Y
 2F2 CDR-H3 (residues 99 to 107): W D R L Y A M D Y


4/23

FIGURE 4

	Tissue origin	Source	CD70 expression*
CA46 Ramos	American Burkitt's lymphoma American Burkitt's lymphoma	ATCC ATCC	3.3 3.2
DAUDI HS-Sultan Raj	African Burkitt's lymphoma African Burkitt's lymphoma African Burkitt's lymphoma	ATCC ATCC ATCC	2.6 5.5 5.4
MC116 Toledo WSU-AHL RL HT 28SR DB MHH-PREB-1 U-698-M SU-DHL-4	Undifferentiated B lymphoma Diffused large cell B lymphoma Histocytic B lymphoma B lymphoblastic lymphoma Diffused mixed B lymphoma Large cell lymphoblastic lymphoma Large cell lymphoblastic lymphoma B lymphoblastic lymphoma B lymphoblastic lymphoma Diffused histiocytic B lymphoma	ATCC ATCC ATCC ATCC ATCC ATCC ATCC DSMZ DSMZ DSMZ	12.2 8.0 11.0 1.3 1.8 1.9 1.4 12.2 3.7 1.1
K1092 L428	Hodgkin's disease Hodgkin's disease	DSMZ DSMZ	25.0 12.9
CESS M-9	B lymphoblastoid cell line B lymphoblastoid cell line	ATCC ATCC	20.7 14.8
Jurkat	Acute T leukemia	ATCC	1.1

* Ratio between the mean log fluorescence intensities for anti-CD70 binding and control IgG binding


5/23

FIGURE 5

		IC50 (μg/ml)	
		1F8-vcMMAE	1F8-vcATP
CA46	American Burkitt's lymphoma	>2	1.4
Ramos	American Burkitt's lymphoma	1.8	>2
DAU1	African Burkitt's lymphoma	1.5	0.5
HS-Sultan	African Burkitt's lymphoma	>2	0.37
Re3	African Burkitt's lymphoma	1.8	>2
MC118	Unfractionated B lymphoma	0.68	0.008
Tob1a	Diffuse large cell B lymphoma	>5	>5
WSU-NHL	Histocytic B lymphoma	0.7	0.09
KOH2	Hodgkin's disease	0.03	0.008
L428	Hodgkin's disease	>2	0.045
CESS	B lymphoblastic cell line	0.017	0.01
Jurkat	T	>2	>2

6/23

FIGURE 6



7/23


FIGURE 7

Sample	Pathology	Age	Sex	CD70 (T.N) ^a
RCC211	clear cell adenocarcinoma	67	M	2.6
RCC213	clear cell adenocarcinoma	74	M	133.8
RCC214	clear cell adenocarcinoma	34	M	92.7
RCC215	clear cell adenocarcinoma	47	M	1.7
RCC216	renal cell carcinoma	57	M	13.8
RCC226	renal cell carcinoma	85	F	0.8
RCC227	renal cell carcinoma	62	F	4.0
RCC306	renal cell carcinoma	49	F	26.0

^aTumor:Normal


8/23

FIGURE 8A




9/23

FIGURE 8B




10/23

FIGURE 9A




11/23

FIGURE 9B



12/23


FIGURE 10

13/23

FIGURE 11

	Proliferation inhibition (IC50, ng/ml)			Cytotoxicity (IC50, ng/ml)	
	1F6-vcMMAE	1F6-vcMMAF	1F6-vcAPP	1F6-vcMMAF	1F6-vcAPP
CalR-1	16	6	9	24	14
CalR-2	10	1	2	3	3
786-O	>5000	6	9	12	15
769-P	2833	24	152	311	1238
SK-RC-6	>5000	ND	12	ND	ND
SK-RC-7	>5000	ND	73	ND	ND
ACHN	>5080	28	247	696	800
CAL54	>5000	11	17	21	240
A498	>5000	2	5	2	2
Hs 835.T	3833	ND	1167	ND	ND
RPTEC	1325	ND	325	>10000	>10000
HRCE	>5000	ND	1150	>10000	>10000


14/23

FIGURE 12


15/23

FIGURE 13

Clone C1A




16/23


FIGURE 14

17/23

FIGURE 15






18/23

FIGURE 10


19/23

FIGURE 17A



20/23


FIGURE 17B



Day 5 after culture initiation


21/23

FIGURE 18




22/23

FIGURE 19



23/23

FIGURE 20



18891-3PC.ST25.txt
SEQUENCE LISTING

<110> Law, Che-Leung
Scholar, Nathalie

<120> ANTI-CD70 ANTIBODIES AND ANTIBODY-DRUG CONJUGATES AND THEIR USE FOR THE
TREATMENT OF CANCER AND IMMUNE DISORDERS

<130> 18891-3PC

<140> PCT/US04/

<141> 2004-02-20

<150> US 60/449,055

<151> 2003-02-20

<160> 47

<170> PatentIn version 3.1

<210> 1

<211> 411

<212> DNA

<213> mouse

<400> 1
atggcttggg tgtggacctt gctattcctg atggcagctg cccaaagtgc ccaagcacag 60
atccagttgg tgcagtctgg acctgaggtg aagaagcctg gagagacagt caagatctcc 120
tgcaaggctt ctgggtatac cttcacaac tatggaatga actgggtgaa gcaggctcca 180
ggaaagggtt taaagtggat gggctggata aacacctaca ctggagagcc aacatatgct 240
gatgccttca agggacgggt tgccttctct ttggaaacct ctgccagcac tgcctatttg 300
cagatcaaca acctcaaaaa tgaggacacg gctacatatt tctgtgcaag agactacggc 360
gactatggta tggactactg gggtaagga acctcagtca ccgtctcctc a 411

<210> 2

<211> 137

18891-3PC.ST25.txt

<212> PRT

<213> mouse

<400> 2

Met Ala Trp Val Trp Thr Leu Leu Phe Leu Met Ala Ala Ala Gln Ser
1 5 10 15

Ala Gly Ala Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Val Lys Lys
20 25 30

Pro Gly Glu Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35 40 45

Thr Asn Tyr Gly Met Asn Trp Val Lys Gln Ala Pro Gly Lys Gly Leu
50 55 60

Lys Trp Met Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala
65 70 75 80

Asp Ala Phe Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser
85 90 95

Thr Ala Tyr Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr
100 105 110

Tyr Phe Cys Ala Arg Asp Tyr Gly Asp Tyr Gly Met Asp Tyr Trp Gly
115 120 125

Gln Gly Thr Ser Val Thr Val Ser Ser
130 135

<210> 3

<211> 57

<212> DNA

<213> mouse

<400> 3

atggcttggtggtgtggaccttgctattcctgatggcagctgcccaaagtgc ccaagca 57

<210> 4

<211> 19

<212> PRT

<213> mouse

18891-3PC.ST25.txt

<400> 4

Met Ala Trp Val Trp Thr Leu Leu Phe Leu Met Ala Ala Ala Gln Ser
1 5 10 15

Ala Gly Ala

<210> 5

<211> 30

<212> DNA

<213> mouse

<400> 5

gggtatacct tcacaaacta tggaatgaac

30

<210> 6

<211> 10

<212> PRT

<213> mouse

<400> 6

Gly Tyr Thr Phe Thr Asn Tyr Gly Met Asn
1 5 10

<210> 7

<211> 51

<212> DNA

<213> mouse

<400> 7

tggataaaca cctacactgg agagccaaca tatgctgatg ccttcaaggg a

51

<210> 8

<211> 17

<212> PRT

<213> mouse

<400> 8

Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Asp Ala Phe Lys
Page 3

18891-3PC.ST25.txt

1 5 10 15

Gly

<210> 9
<211> 27
<212> DNA
<213> mouse

<400> 9
gactacggcg actatggtat ggactac 27

<210> 10
<211> 9
<212> PRT
<213> mouse

<400> 10
Asp Tyr Gly Asp Tyr Gly Met Asp Tyr
1 5

<210> 11
<211> 396
<212> DNA
<213> mouse

<400> 11
atggagacag acacactcct gttatgggta ctgctgctct gggttccagg ttccactggt 60
gacattgtgc tgacacagtc tcctgcttcc ttagctgtat ctctggggca gagggccacc 120
atctcatgca gggccagcaa aagtgtcagt acatctggct atagttttat gcactggtat 180
caacagaaac caggacagcc acccaaaactc ctcattctatc ttgcatccaa cctagaatct 240
ggggtccctg ccagggttcag tggcagtggg tctgggacag acttcaccct caacatccat 300
cctgtggagg aggaggatgc tgcaacctat tactgtcagc acagtaggga gggtccgtgg 360
acgttcggtg gaggcaccaa gctggaaatc aaacgg 396

<210> 12
<211> 132
<212> PRT

18891-3PC.ST25.txt

<213> mouse

<400> 12

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1 5 10 15

Gly Ser Thr Gly Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala
 20 25 30

Val Ser Leu Gly Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Lys Ser
 35 40 45

Val Ser Thr Ser Gly Tyr Ser Phe Met His Trp Tyr Gln Gln Lys Pro
 50 55 60

Gly Gln Pro Pro Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser
 65 70 75 80

Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 85 90 95

Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys
 100 105 110

Gln His Ser Arg Glu Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu
 115 120 125

Glu Ile Lys Arg
 130

<210> 13

<211> 60

<212> DNA

<213> mouse

<400> 13

atggagacag acacactcct gttatgggta ctgctgctct gggttccagg ttccactggg 60

<210> 14

<211> 20

<212> PRT

<213> mouse

<400> 14

18891-3PC.ST25.txt

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
1 5 10 15

Gly Ser Thr Gly
20

<210> 15

<211> 45

<212> DNA

<213> mouse

<400> 15
agggccagca aaagtgtcag tacatctggc tatagtttta tgcac

45

<210> 16

<211> 15

<212> PRT

<213> mouse

<400> 16

Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser Phe Met His
1 5 10 15

<210> 17

<211> 21

<212> DNA

<213> mouse

<400> 17
cttgcaccca acctagaatc t

21

<210> 18

<211> 7

<212> PRT

<213> mouse

<400> 18

Leu Ala Ser Asn Leu Glu Ser
1 5

18891-3PC.ST25.txt

<210> 19
 <211> 27
 <212> DNA
 <213> mouse

<400> 19
 cagcacagta gggaggttcc gtggacg

27

<210> 20
 <211> 9
 <212> PRT
 <213> mouse

<400> 20
 Gln His Ser Arg Glu Val Pro Trp Thr
 1 5

<210> 21
 <211> 411
 <212> DNA
 <213> mouse

<400> 21
 atggaatgga cctgggtctt tctcttcctc ctgccagtaa ctgcagatgt ccaatcccag 60
 gttcagctgc aacagtctgg aactgagctg atgacgcctg gggcctcagt gacgatgtcc 120
 tgcaagactt ctggctacac attcagtacc tactggatag agtgggtaaa acagaggcct 180
 ggacatggcc ttgagtggat tggagaaatt ttacctggaa gtgggttatac tgactacaat 240
 gagaagttca aggccaaggc cacattcact gcagatacat cctccaacac agcctacatg 300
 caactcagca gcctggcatc tgaggactct gccgtctatt actgtgcaag atgggatagg 360
 ctctatgcta tggactactg gggtaagga acctcagtca ccgtctcctc a 411

<210> 22
 <211> 137
 <212> PRT
 <213> mouse

<400> 22

18891-3PC.ST25.txt

Met Glu Trp Thr Trp Val Phe Leu Phe Leu Leu Ser Val Thr Ala Asp
 1 5 10 15

Val Gln Ser Gln Val Gln Leu Gln Gln Ser Gly Thr Glu Leu Met Thr
 20 25 30

Pro Gly Ala Ser Val Thr Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe
 35 40 45

Ser Thr Tyr Trp Ile Glu Trp Val Lys Gln Arg Pro Gly His Gly Leu
 50 55 60

Glu Trp Ile Gly Glu Ile Leu Gly Pro Ser Gly Tyr Thr Asp Tyr Asn
 65 70 75 80

Glu Lys Phe Lys Ala Lys Ala Thr Phe Thr Ala Asp Thr Ser Ser Asn
 85 90 95

Thr Ala Tyr Met Gln Leu Ser Ser Leu Ala Ser Glu Asp Ser Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Trp Asp Arg Leu Tyr Ala Met Asp Tyr Trp Gly
 115 120 125

Gly Gly Thr Ser Val Thr Val Ser Ser
 130 135

<210> 23

<211> 57

<212> DNA

<213> mouse

<400> 23

atggaatgga cctgggtctt tctcttcctc ctgctagtaa ctgcagatgt ccaatcc

57

<210> 24

<211> 19

<212> PRT

<213> mouse

<400> 24

Met Glu Trp Thr Trp Val Phe Leu Phe Leu Leu Ser Val Thr Ala Asp
 1 5 10 15

Val Gln Ser

18891-3PC.ST25.txt

<210> 25

<211> 30

<212> DNA

<213> mouse

<400> 25

ggctacacat tcagtaccta ctggatagag

30

<210> 26

<211> 10

<212> PRT

<213> mouse

<400> 26

Gly Tyr Thr Phe Ser Thr Tyr Trp Ile Glu
1 5 10

<210> 27

<211> 51

<212> DNA

<213> mouse

<400> 27

gaaattttac ctggaagtgg ttatactgac tacaatgaga agttcaaggc c

51

<210> 28

<211> 17

<212> PRT

<213> mouse

<400> 28

Glu Ile Leu Gly Pro Ser Gly Tyr Thr Asp Tyr Asn Glu Lys Phe Lys
1 5 10 15

Ala

<210> 29

18891-3PC.ST25.txt

<211> 27

<212> DNA

<213> mouse

<400> 29

tgggataggc tctatgctat ggactac

27

<210> 30

<211> 9

<212> PRT

<213> mouse

<400> 30

Trp Asp Arg Leu Tyr Ala Met Asp Tyr
1 5

<210> 31

<211> 396

<212> DNA

<213> mouse

<400> 31

atggagacag acacactcct gttatgggta ctgctgctct gggttccagg ttccactggt	60
gacattgtgc tgacacagtc tcctgcttcc ttaactgtat ctctggggca gaagaccacc	120
atctcatgca gggccagcaa gagtgtcagt acatctggct atagttttat gcactggtac	180
caactgaaac caggacagtc acccaaactc ctcattctatc ttgcgtccaa cctaccatct	240
ggggtccctg ccaggttcag tggcagtggg tctgggacag acttcaccct caaaatccat	300
cctgtggagg aggaggatgc tgcaacctat tactgtcagc acagtaggga gattccgtac	360
acgttcggag gggggaccaa gctggaaata acacgg	396

<210> 32

<211> 132

<212> PRT

<213> mouse

<400> 32

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro

18891-3PC.ST25.txt

1 5 10 15

Gly Ser Thr Gly Asp Ile Val Leu Thr Gln Ser pro Ala Ser Leu Thr
 20 25 30

Val Ser Leu Gly Gln Lys Thr Thr Ile Ser Cys Arg Ala Ser Lys Ser
 35 40 45

Val Ser Thr Ser Gly Tyr Ser Phe Met His Trp Tyr Gln Leu Lys Pro
 50 55 60

Gly Gln Ser Pro Lys Leu Leu Ile Tyr Leu Ala Ser Asp Leu Pro Ser
 65 70 75 80

Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 85 90 95

Leu Lys Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys
 100 105 110

Gln His Ser Arg Glu Ile Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu
 115 120 125

Glu Ile Thr Arg
 130

<210> 33

<211> 60

<212> DNA

<213> mouse

<400> 33

atggagacag acacactcct gttatgggta ctgctgctct gggttccagg ttccactggg 60

<210> 34

<211> 20

<212> PRT

<213> mouse

<400> 34

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1 5 10 15Gly Ser Thr Gly
 20

18891-3PC.ST25.txt

<210> 35

<211> 45

<212> DNA

<213> mouse

<400> 35

agggccagca agagtgtcag tacatctggc tatagtttta tgcac

45

<210> 36

<211> 15

<212> PRT

<213> mouse

<400> 36

Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser Phe Met His
1 5 10 15

<210> 37

<211> 21

<212> DNA

<213> mouse

<400> 37

cttgcgtcca acctaccatc t

21

<210> 38

<211> 7

<212> PRT

<213> mouse

<400> 38

Leu Ala Ser Asn Leu Pro Ser
1 5

<210> 39

<211> 28

<212> DNA

<213> mouse

18891-3PC.ST25.txt

<400> 39
cagcacagta gggagattcc gtacacgt

28

<210> 40

<211> 9

<212> PRT

<213> mouse

<400> 40

Gln His Ser Arg Glu Ile Pro Tyr Thr
1 5

<210> 41

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 41
cttcacttg acattgatgt ctttg

25

<210> 42

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 42
caggtcactg tcactggctc ag

22

<210> 43

<211> 37

<212> DNA

<213> Artificial Sequence

18891-3PC.ST25.txt

<220>

<223> Primer

<400> 43

gtcgatgagc tctagaattc gtgccccccc ccccccc

37

<210> 44

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 44

cgtcatgtcg acggatccaa gcttcaagaa gcacacgact gaggcac

47

<210> 45

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 45

cgtcatgtcg acggatccaa gcttgtcacc atggagttag tttgggc

47

<210> 46

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer

<400> 46

ccactgctgc tgattag

17

<210> 47

<211> 18

<212> DNA

18891-3PC.ST25.txt

<213> Artificial Sequence

<220>

<223> Primer

<400> 47
caatgccttc tcttgtcc

18

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.